

Specimen Label



Gallery[®]

75 Dry Flowable

Specialty Herbicide

®Trademark of Dow AgroSciences LLC

A preemergence herbicide for control of certain broadleaf weeds in:

- Established Turfgrass
- Landscape Ornamentals
- Container Grown Ornamentals
- Field Grown Ornamentals
- Groundcovers/Perennials
- Non-Cropland
- Ornamental Bulbs
- Non-Bearing Fruit and Nut Trees and Non-Bearing Vineyards
- Christmas Tree/Conifer Plantations

Active Ingredient:

isoxaben: N-[3-(1-ethyl-1-methylpropyl)- 5-isoxazolyl]-

2,6-dimethoxybenzamide and isomers 75%

Other Ingredients 25%

Total 100%

Contains 0.75 lb active ingredient per pound.

U.S. Patent Nos. 5,086,184 and 4,636,243

EPA Reg. No. 62719-145

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Eye Irritation • Harmful If Inhaled

Avoid ingestion, breathing dust or spray mist, and contact with skin, eyes, or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift may result in reduced germination or emergence of non-target plants adjacent to treated area.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: When this product is applied to turf and ornamental plantings in landscape settings and non-cropland areas, do not allow entry into treated areas until sprays have dried unless wearing coveralls, waterproof gloves, and shoes plus socks.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container. Do not store in direct sunlight. Do not store at temperatures above 120°F. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Storage and Disposal (Cont.)

Nonrefillable rigid containers 5 gallons or less:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Nonrefillable nonrigid containers:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refillable rigid containers larger than 5 gal:

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Nonrefillable rigid containers larger than 5 gal:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

General Information

Gallery® 75 Dry Flowable specialty herbicide is a preemergence product for control of certain broadleaf weeds in established turfgrass, landscape ornamentals, container grown ornamentals, field grown ornamentals, groundcovers/perennials, ornamental bulbs, non-bearing fruit and nut trees and non-bearing vineyards, Christmas tree/conifer plantations and non-cropland areas. Apply Gallery 75 Dry Flowable in late summer to early fall, in early spring, or anytime prior to germination of target weeds, or immediately after cultivation.

Use Precautions and Restrictions

Gallery 75 Dry Flowable controls weeds germinating from seed. Gallery 75 Dry Flowable does not control established weeds, or weeds growing from stolons, rhizomes, or root pieces. Existing weeds should be controlled by cultivation or with postemergence herbicides. Weed residues, prunings, and trash should be removed or thoroughly mixed into the soil prior to application. Soil in non-turf areas should be in good condition and free of clods at the time of application. Gallery 75 Dry Flowable is stable on the soil surface for up to 21 days, but must be incorporated by moisture to be effective. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation after application is necessary to activate Gallery 75 Dry Flowable. If Gallery 75 Dry Flowable is not activated by rainfall or irrigation within 21 days after application, erratic weed control may result. In non-turfgrass areas, if weeds emerge due to lack of rainfall or irrigation, shallow cultivation to a depth of 1 to 2 inches will incorporate the herbicide and destroy existing weeds.

Treatment of Turfgrass or Ornamental Species Not Listed on the Label for Gallery 75 Dry Flowable:

Although this label contains a large number of ornamental species, it is not possible to include all of the ornamental plants that may be encountered in nursery or landscape settings. Users who wish to use Gallery 75 Dry Flowable on a plant species not listed on this label may determine the suitability for such use by treating a small area or small number of plants at a recommended rate. Prior to treatment of larger areas, the treated area/plants should be observed for any sign of herbicidal injury during 30 to 60 days of typical growing conditions. The user assumes the responsibility for any plant damage or other liability resulting from use of Gallery 75 Dry Flowable on species not recommended on this label.

In Arizona: The state of Arizona has not approved Gallery 75 Dry Flowable for use on plants grown for commercial production such as: turf sod farms, Christmas tree plantations, ornamentals and nursery stock grown for resale, and non-bearing fruit and nut trees and non-bearing vineyards.

Chemigation: Do not apply Gallery 75 Dry Flowable through any type of irrigation system.

Application Directions

Apply Gallery 75 Dry Flowable with a properly calibrated low-pressure herbicide sprayer that provides uniform spray distribution. Nozzle screens should be no finer than 50 mesh (50 mesh is finer than 16 mesh). In-line screens and strainers should be no finer than 16 mesh. Apply Gallery 75 Dry Flowable in 10 gallons or more of water carrier per acre. As the spray volume decreases, the importance of accurate calibration and uniform application increases. Take precautions to avoid spray drift when applying Gallery 75 Dry Flowable. Drift may result in reduced germination or emergence of non-target plants adjacent to the treated area. Maintain agitation from mixing through application. Avoid boom overlaps that will increase rates above those recommended. Calibrate application equipment prior to use, according to manufacturer's directions. Check calibration frequently to be sure equipment is working properly and distributing spray uniformly.

Mixing Directions

Gallery 75 Dry Flowable - Alone

Check to be sure spray equipment is clean and not contaminated with other herbicides. Using clean water, fill the tank to 1/2 of the final volume required and start agitation. Slowly add the required quantity of Gallery 75 Dry Flowable to the spray tank, continue agitation and complete filling the tank. Maintain agitation during filling and throughout application. Agitation should be sufficient to create a rippling or rolling action on the liquid surface. Sparger pipe agitation generally provides the best agitation. In-line screens and strainers should be no finer than 16 mesh.

If spraying and agitation is stopped, Gallery 75 Dry Flowable may settle to the bottom of the spray tank. If settling occurs, material must be re-suspended before continuing spray application. Clean the spray tank, lines and screens thoroughly after use.

Application Rate Conversion Table for Gallery 75 Dry Flowable

lb per acre	oz per 1000 sq ft	grams per 1000 sq ft
0.66	0.25	7.0
1.00	0.38	10.5
1.33	0.50	14.0

Repeat applications at 1 lb or more per acre of Gallery 75 Dry Flowable should not be made sooner than 60 days after a previous application of Gallery 75 Dry Flowable. Do not apply more than a total of 4 lb of Gallery 75 Dry Flowable per acre within a 12-month period.

Note: The container cap for Gallery 75 Dry Flowable may be used to measure the amount of product required for small areas. When filled to the level of the inner lip, the cap contains approximately 1/2 oz, which is the amount of Gallery 75 Dry Flowable needed to treat an area of 1000 sq ft at a rate equivalent to 1.33 lb per acre.



When filled to this level, container cap contains 1/2 oz (14 grams) of Gallery 75 Dry Flowable.

Gallery 75 Dry Flowable - Tank Mix

Gallery 75 Dry Flowable may be applied in tank mix combination with labeled rates of other products provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; (2) tank mixing with Gallery 75 Dry Flowable is not prohibited by the label of the tank mix product; and (3) the tank mix combination is compatible as determined by a "jar test" described in the Tank Mix Compatibility Testing section below.

Add Gallery 75 Dry Flowable to the spray tank as described above and fill the spray tank to 3/4 of the final volume required. Add other formulations in this order: (1) other dry flowables, (2) wettable powders, (3) aqueous suspensions, (4) flowables, (5) liquids, (6) solutions and emulsifiable or liquid concentrates. Allow two to three minutes between the addition of each product and agitate continuously until the product(s) is completely dispersed in water. Maintain agitation during filling and through application. If a build up of materials is observed on the walls of the spray tank, wash the tank with soapy water between fillings, rinse and then continue the spraying operation. Follow label directions for each material added to the tank.

Tank Mix Compatibility Testing: A jar test is recommended prior to tank mixing to ensure compatibility of Gallery 75 Dry Flowable and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

Weeds Controlled or Suppressed

Weeds controlled when applied at 0.66 lb per acre (0.25 oz or 7 grams/1000 sq ft):

Common Name
aster, slender
bursage, annual
burweed, lawn
celery, wild
chickweed, common
clover, white
cudweed, purple
fiddleneck, coast
filaree, redstem
fleabane, blackleaved
fleabane, dwarf
groundcherry, lanceleaf
henbit
horseweed
knotweed, prostrate
lambsquarters, common
mallow, little
mustard, Indian
mustard, wild

Scientific Name
Aster exilis
Ambrosia acanthicarpa
Soliva pterosperma
Apium leptophyllum
Stellaria media
Trifolium repens
Gnaphalium purpureum
Amsinckia intermedia
Erodium cicutarium
Conyza bonariensis
Conyza ramosissima
Physalis lanceifolia
Lamium amplexicaule
Conyza canadensis
Polygonum aviculare
Chenopodium album
Malva parviflora
Brassica juncea
Sinapis arvensis

nightshade, black
pepperweed, Virginia
pigweed
pineappleweed
plantain, slender
purslane, common
radish, wild
ragweed, common
rocket, London
rockpurslane, desert
shepherdspurse
sibara
smartweed, Pennsylvania
sowthistle, annual
speedwell, purslane
telegraphplant
thistle, Russian

Solanum nigrum
Lepidium virginicum
Amaranthus spp.
Matricaria matricarioides
Plantago elongata
Portulaca oleracea
Raphanus raphanistrum
Ambrosia artemisiifolia
Sisymbrium irio
Calandrinia ciliata
Capsella bursa-pastoris
Sibara virginica
Polygonum pensylvanicum
Sonchus oleraceus
Veronica peregrina
Heterotheca grandiflora
Salsola iberica

Weeds controlled when applied at 1 lb per acre (0.38 oz or 10.5 grams/1000 sq ft):

Common Name
aster, heath
bittercress
bittercress, hairy
brassbuttons, southern
carrot, wild
chamberbitter
chickweed, mouseear
dandelion
eclipta
galinsoga, hairy
geranium, carolina
ladysthumb
lettuce, prickly
mallow, dwarf
marestail (see horseweed)
mayweed
morningglory, ivyleaf
mustard, black
pennywort
phyllanthus, long-stalk
plantain, bracted
plantain, broadleaf
plantain, buckhorn
pokeweed, common
rockpurslane, redmaids
sida, prickly
sorrell, red
speedwell, thymeleaf
spurge, hyssop
spurge, spotted
sweetclover, yellow
tansymustard, green

woodsorrel, yellow

Scientific Name
Aster ericoides
Cardamine oligosperma
Cardamine hirsuta
Cotula australis
Daucus carota
Phyllanthus urinaria
Cerastium vulgatum
Taraxacum officinale
Eclipta prostrata
Galinsoga ciliata
Geranium carolinianum
Polygonum persicaria
Lactuca serriola
Malva rotundifolia

Anthemis cotula
Ipomoea hederacea
Brassica nigra
Hydrocotyle spp.
Phyllanthus tenellus
Plantago aristata
Plantago major
Plantago lanceolata
Phytolacca americana
Calandrinia ciliata var. *menziesii*
Sida spinosa
Rumex acetosella
Veronica serpyllifolia
Euphorbia hyssopifolia
Euphorbia maculata
Melilotus officinalis
Descurainia pinnata
ssp. *brachycarpa*
Oxalis stricta

Weeds controlled when applied at 1.33 lb per acre (0.5 oz or 14 grams/1000 sq ft):

Common Name	Scientific Name
burclover, California	<i>Medicago polymorpha</i>
dogfennel	<i>Eupatorium capillifolium</i>
eveningprimrose	<i>Oenothera</i> spp.
fescue, rattail	<i>Vulpia myuros</i>
filaree, whitestem	<i>Erodium moschatum</i>
goosefoot, nettleleaf	<i>Chenopodium murale</i>
groundsel, common	<i>Senecio vulgaris</i>
jimsonweed	<i>Datura stramonium</i>
knotweed, silversheath	<i>Polygonum argyrocoleon</i>
kochia	<i>Kochia scoparia</i>
medic, black	<i>Medicago lupulina</i>
mullein, turkey	<i>Eremocarpus setigerus</i>
nettle, burning	<i>Urtica urens</i>
oxtongue, bristly	<i>Picris echioides</i>
pimpernel, scarlet	<i>Anagallis arvensis</i>
sowthistle, spiny	<i>Sonchus asper</i>
spurge, petty	<i>Euphorbia peplus</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
sunflower	<i>Helianthus</i> spp.
swinecress	<i>Coronopus didymus</i>
thistle, musk	<i>Carduus nutans</i>
willoweed, panicle	<i>Epilobium paniculatum</i>
woodsorrel, creeping	<i>Oxalis corniculata</i>

Weeds partially controlled or suppressed when applied at 1.33 lb per acre (0.5 oz or 14 grams/1000 sq ft):

Common Name	Scientific Name
bindweed, field	<i>Convolvulus arvensis</i>
carpetweed	<i>Mollugo verticillata</i>
dock, curly	<i>Rumex crispus</i>
mallow, Venice	<i>Hibiscus trionum</i>
milkweed, honeyvine	<i>Ampelamus albidus</i>
morningglory, tall	<i>Ipomoea purpurea</i>
pusley, Florida	<i>Richardia scabra</i>

Uses

Established Turfgrass

Gallery 75 Dry Flowable is recommended as a preemergence treatment for control of certain broadleaf weeds in established cool season and warm season turfgrass.

Apply Gallery 75 Dry Flowable anytime prior to germination of target weeds.

Note: Refer to the General Information section of this label for Use Precautions and Restrictions and information on mixing and application, application rates, and weeds controlled prior to using this product.

Use Precautions:

Apply Gallery 75 Dry Flowable to newly seeded turfgrass (including overseeded turfgrass) **only** after seedlings are established (three leaf stage and tillering) and well rooted. Do not overseed established turfgrass sooner than 60 days following an application of Gallery 75 Dry Flowable.

- Do not apply Gallery 75 Dry Flowable to golf course putting greens or tees.
- Do not apply Gallery 75 Dry Flowable to dichondra.
- Do not apply Gallery 75 Dry Flowable to turfgrass grown for seed.

Tank Mixing

Gallery 75 Dry Flowable may be tank mixed with Dimension® herbicide and applied as a preemergence treatment to **warm season turfgrass only** to broaden the spectrum of annual grass and broadleaf weed control. Gallery 75 Dry Flowable may also be applied as a separate treatment to supplement the effectiveness of Balan® 2.5G or Team® 2G herbicides in cool and warm season turfgrass. Gallery 75 Dry Flowable may be tank mixed with postemergence broadleaf herbicides registered for use on established turfgrass to control existing broadleaf weeds to provide residual preemergence broadleaf weed control. Applied as directed, Gallery 75 Dry Flowable in tank mix with other products registered for use on turfgrass will provide control of susceptible weed species listed on the respective labels. When using Gallery 75 Dry Flowable in tank mix combinations with other products, read and follow all applicable use directions, precautions, and limitations on the respective product labels. Refer to instructions for Gallery 75 Dry Flowable in Tank Mix in the Mixing Directions section.

Gallery 75 Dry Flowable is recommended for use on the following turfgrass species:

Common Name	Scientific Name
Established Cool Season Turfgrass	
bentgrass, creeping	<i>Agrostis stolonifera</i>
bentgrass, colonial	<i>Agrostis tenuis</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, chewing	<i>Festuca rubra</i> var. <i>commutata</i>
fescue, creeping red	<i>Festuca rubra</i>
fescue, sheeps	<i>Festuca ovina</i>
fescue, tall	<i>Festuca arundinaceae</i>
ryegrass, perennial	<i>Lolium perenne</i>
Established Warm Season Turfgrass¹	
bahiagrass	<i>Paspalum notatum</i>
bermudagrass	<i>Cynodon dactylon</i>
buffalograss	<i>Buchloe dactyloides</i>
centipedegrass	<i>Eremochloa ophiuroides</i>
fescue, tall (growing in warm season areas)	<i>Festuca arundinaceae</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
zoysiagrass	<i>Zoysia japonica</i>
zoysiagrass	<i>Zoysia tenuifolia</i>

¹ **Sprigged Warm Season Turfgrass:** Gallery 75 Dry Flowable is recommended post-sprigging as a preemergence treatment for control of certain broadleaf weeds in warm season turfgrass. Apply anytime after sprigging in the following turfgrass species: bermudagrass, bahiagrass, St. Augustinegrass, centipedegrass and buffalograss. Do not apply more than 1 lb of Gallery 75 Dry Flowable per acre during the establishment phase for newly sprigged warm season turfgrass. Do not apply Gallery 75 Dry Flowable to varieties of dwarf-type bermudagrass or to any turfgrass species being sprigged on golf course tees or greens.

Ornamental Plantings and Non-Bearing Fruit and Nut Trees and Non-Bearing Vineyards

Gallery 75 Dry Flowable is recommended as a preemergence treatment for control of certain broadleaf weeds in landscape ornamentals, container grown ornamentals, field grown ornamentals, groundcovers/perennials, and non-bearing fruit and nut trees and non-bearing vineyards.

Apply Gallery 75 Dry Flowable anytime prior to germination of target weeds, or immediately after cultivation.

Note: Refer to the General Information section of this label for Use Precautions and Restrictions and information on mixing and application, application rates, and weeds controlled prior to using this product.

Tank Mixing

Gallery 75 Dry Flowable may be tank mixed with Roundup or other postemergence herbicides registered for control of existing unwanted vegetation in ornamental plantings and non-bearing fruit and nut trees and non-bearing vineyards and to provide residual preemergence broadleaf weed control. Gallery 75 Dry Flowable may also be tank mixed with Dimension and applied preemergence to provide broad spectrum control of annual grasses and broadleaf weeds in ornamental areas and non-bearing fruit and nut trees and non-bearing vineyards. Applied as directed, tank mixes of Gallery 75 Dry Flowable will provide control of susceptible weed species listed on the respective labels. When using Gallery 75 Dry Flowable in tank mix combination with other products, read and follow all applicable use directions, precautions, and limitations on the respective product labels. Refer to instructions for Gallery 75 Dry Flowable in Tank Mix in the Mixing Directions section.

Note: Do not apply sprays containing Roundup over the top of ornamental plants. Extreme care must be exercised to prevent contact of sprays containing Roundup with foliage or stems of turfgrass, trees, shrubs, or other desirable vegetation since severe damage or death may result. If spraying Roundup in areas adjacent to desirable plants, use a shield to prevent spray from contacting foliage or stems of desirable plants.

Use Precautions:

To avoid possible plant injury, **do not** apply Gallery 75 Dry Flowable to:

- Nursery, forest, or Christmas tree: seedling beds, cutting beds, or transplant beds
- Unrooted liners or cuttings that have been planted in pots for the first time
- Pots less than six inches wide
- Groundcovers until they are established and well rooted
- Bedding plants or areas where bedding plants will be planted or transplanted within one year after application

Do not apply Gallery 75 Dry Flowable to newly transplanted ornamentals, nursery stock, groundcovers, ornamental bulbs, non-bearing fruit and nut trees or non-bearing vineyards until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or plant injury may occur.

Applications of Gallery 75 Dry Flowable over the top of plants with newly forming buds may cause injury. Possible plant injury may be avoided by application as a directed spray to the soil surface beneath ornamental plants.

When planting into a site treated with Gallery 75 Dry Flowable in the past 8 months, use untreated soil as fill around roots when replacing plants or injury may occur.

Note: Injury to certain ornamental plants has been observed following application of Gallery 75 Dry Flowable. To avoid plant injury, Gallery 75 Dry Flowable is not recommended for weed control in the following ornamental plant species:

<i>Ajuga</i> spp.	bugle
<i>Brassica</i> spp.	mustard
<i>Echinacea purpurea</i>	purple coneflower
<i>Euonymus alatus 'compacta'</i>	dwarf burning bush
<i>Euphorbia</i> spp.	spurge
<i>Hydrangea</i> spp.	hydrangea
<i>Iberis</i> spp.	candytuft
<i>Juniperus horizontalis 'Prince of Wales'</i>	prince of wales juniper
<i>Melaleuca quinquenervia</i>	cajeput tree
<i>Rhododendron caroliniaum</i>	carolina rhododendron
<i>Rhododendron catawbiense 'Roseum elegans'</i>	roseum elegans rhododendron
<i>Sedum</i> spp.	stonecrop
<i>Yucca recurvifolia</i>	green yucca

Gallery 75 Dry Flowable may be used in the culture of the following established plant species: (**Note:** Limitations on recommended treatment methods)

Trees

Recommended Treatment Method:

C = Container Grown

F = Field Grown

Scientific Name	Common Name	
<i>Abies balsamea</i>	fir, balsam	C, F
<i>Abies concolor</i>	fir, white	F
<i>Abutilon hybridum</i>	albus-flowering maple	C, F
	luteus-flowering maple	C, F
	roseus-flowering maple	C, F
	tangerine-flowering maple	C, F
	vesuvius red-flowering maple	F
<i>Acer ginnala</i>	flame maple	F
<i>Acer rubrum</i>	red maple	F
	red sunset maple	F
<i>Acer saccharinum</i>	silver maple	C, F
<i>Alsophila australis</i>	Australian tree fern	C, F
<i>Archontophoenix cunninghamiana</i>	king palm	C, F
<i>Areacastrum romanzoffianum</i>	queen palm	C, F
<i>Araucaria heterophylla</i>	Norfolk island pine	C, F
<i>Bauhinia galpinii</i>	red bauhinia	C, F
<i>Betula nigra</i>	birch, river	C, F
<i>Betula papyrifera</i>	paper birch	F
<i>Brachychiton populneus</i>	bottle tree	C, F
<i>Bucida buceras</i>	black olive	F
<i>Ceratonia siliqua</i>	carob	F
<i>Cercis canadensis</i>	redbud	C, F
<i>Chamaecyparis obtusa</i>	filicoides-fernspray cypress	F
	gracilis-slender hinoki cypress	F
<i>Chamaecyparis pisifera</i>	sawara-false cypress	F
	squarrosa-moss cypress	F

Trees (Cont.)

Recommended Treatment Method:
C = Container Grown
F = Field Grown

Scientific Name	Common Name	
<i>Chamaedorea cataractarum</i>	cat palm	F
	palm	C, F
<i>Chamaedorea costaricana</i>	palm	C, F
<i>Chamaedorea elegans</i>	parlor palm	C, F
<i>Chamaerops humilis</i>	Mediterranean fan palm	C, F
<i>Cornus florida</i>	cloud nine dogwood	C, F
	dogwood, flowering	C, F
<i>Crataegus viridis</i>	green hawthorn	F
<i>Cryptomeria japonica</i>	cryptomeria, Japanese	C, F
<i>Cupaniopsis anacardioides</i>	carrot wood	F
<i>Cupressus glabra</i>	Arizona cypress	F
<i>Cupressocyparis 'Emerald Isle'</i>	emerald Isle leyland cypress	C, F
<i>Cupressocyparis leylandii</i>	blue leyland cypress	C, F
<i>Cupressus ariz 'Blue pyramid'</i>	blue pyramid cypress	C, F
<i>Cupressus sempervirens</i>	cypress tree	C, F
<i>Cycas revoluta</i>	sago palm	C, F
<i>Elaeagnus angustifolia</i>	Russian olive	C, F
<i>Elaeagnus x ebbengei</i>	gilt edge silverberry	C, F
	'Gilt edge'	
<i>Eucalyptus camaldulensis</i>	red gum eucalyptus	F
<i>Eucalyptus cinerea</i>	eucalyptus, mealy	F
	silver dollar eucalyptus	F
<i>Eucalyptus microtheca</i>	coolibah tree	C, F
<i>Eucalyptus sideroxylon</i>	eucalyptus, red ironbark	F
<i>Fagus sylvatica</i>	European beech	C, F
<i>Ficus benjamina</i>	ficus	C, F
	mini ficus	C, F
<i>Fraxinus udhei</i>	shamel ash	C, F
<i>Ginkgo biloba</i>	ginkgo (maidenhair tree)	F
<i>Gleditsia triacanthos</i>	honey locust	F
	shademaster honey locust	F
<i>Heteromeles arbutiflora</i>	toyon	F
<i>Illicium floridanum</i>	Florida anise-tree	C, F
<i>Juniperus virginiana</i>	redcedar, eastern	C, F
<i>Leptospermum scoparium</i>	tea tree	C, F
<i>Liquidambar styraciflua</i>	sweetgum, American	F
<i>Magnolia grandiflora</i>	magnolia, southern	C, F
<i>Magnolia stellata</i>	royal star magnolia	C, F
<i>Morus alba</i>	white mulberry	F
<i>Musa aluminata</i>	banana	C, F
<i>Oxydendrum arboreum</i>	sourwood	C, F
<i>Picea abies</i>	pendula-weeping Norway spruce	C, F
	repens-spreading Norway spruce	C, F
	spruce, Norway	C, F
<i>Picea glauca</i>	conica-dwarf alberta spruce	C, F
<i>Picea glauca conica</i>	conica-dwarf alberta spruce	F
<i>Picea pungens</i>	dwarf globe blue spruce	C, F
	glauca-Colorado blue spruce	C, F
	hoopsii-hoop's blue spruce	C, F
	koster-koster blue spruce	F
<i>Pinus aristata</i>	bristlecone pine	F
<i>Pinus canariensis</i>	canary Island pine	F
<i>Pinus contorta</i>	shore pine, beach pine	F
<i>Pinus eldarica</i>	eldarica pine	C, F
<i>Pinus leucodermis</i>	Bosnian pine	C, F
<i>Pinus mugo</i>	pumilio- shrubby swiss	C, F
	mountain pine	
<i>Pinus nigra</i>	Austrian black pine	C, F
<i>Pinus radiata</i>	monterey pine	F

<i>Pinus strobus</i>	eastern white pine	C, F
	white pine	C, F
<i>Pinus sylvestris</i>	columnar Scotch pine	C, F
	Scotch pine	C, F
<i>Pinus thunbergiana</i>	Japanese black pine	C, F
<i>Platanus occidentalis</i>	American sycamore	F
<i>Platanus racemosa</i>	Californina sycamore	F
<i>Podocarpus spp.</i>	podocarpus	F
<i>Populus deltoides</i>	cottonwood	F
<i>Prosopis chilensis</i>	Chilean mesquite	C, F
<i>Prunus yedoensis</i>	yoshino flowering cherry	F
<i>Prunus caroliniana</i>	carolina laurel cherry	C, F
<i>Prunus laurocerasus</i>	English laurel	C, F
<i>Quercus ilicifolia</i>	bear oak	F
<i>Quercus palustris</i>	pin oak	F
<i>Quercus phellos</i>	willow oak	C, F
<i>Quercus rubra</i>	red oak	C, F
<i>Quercus virginiana</i>	live oak	C, F
<i>Salix babylonica</i>	babylon weeping willow	F
	corkscrew willow	F
<i>Sequoiadendron giganteum</i>	giant sequoia	F
<i>Swietenia mahogany</i>	mahogany	F
<i>Tabebuia caraiba</i>	yellow tab	F
<i>Trachycarpus fortunei</i>	windmill palm	C, F
<i>Tsuga canadensis</i>	eastern hemlock	C, F
<i>Ulmus parvifolia</i>	Chinese elm	F
<i>Washingtonia robusta</i>	Mexican fan palm	F
<i>Zamia furfuracea</i>	cardboard palm	C, F

Shrubs

Recommended Treatment Method:
C = Container Grown
F = Field Grown

Scientific Name	Common Name	
<i>Abelia grandiflora</i>	edward goucher abelia	C, F
	glossy abelia	C, F
	sunrise variegated abelia	C, F
<i>Acacia abyssinica</i>	abyssinica acacia	C, F
<i>Acacia redolens</i>	acacia, prostrate	C, F
<i>Acacia stenophylla</i>	shoestring acacia	C, F
<i>Acalypha wilkesiana</i>	copper leaf	C, F
<i>Acer ginnala</i>	amur maple	C, F
<i>Acer palmatum</i>	coral bark Japanese maple	C, F
<i>Acer palmatum</i>	dwarf Japanese maple	C, F
<i>Agave americana</i>	century plant	F
<i>Amelanchier alnifolia</i>	saskatoon serviceberry	C, F
<i>Andromeda polifolia</i>	bog rosemary	C, F
<i>Anisodonteia hypomandarum</i>	cape mallow	C, F
<i>Arctostaphylos uva-ursi</i>	bearberry	C, F
<i>Ardisia japonica</i>	chirimen marlberry	C, F
<i>Artemesia lactiflora</i>	white mugwort	C, F
<i>Athyrium nipponicum</i>	Japanese painted fern	C, F
<i>Baccharis pilularis</i>	coyotebush	F
<i>Berberis gladwynensis</i>	william penn barberry	C, F
<i>Berberis mentorensis</i>	mentor barberry	C, F
<i>Berberis thunbergii</i>	aurea-golden Japanese barberry	F
	crimson pygmy barberry	C, F
	rose glow barberry	C, F
	atropurea-redleaf Japanese barberry	F
	cherry bomb	C, F

Shrubs (Cont.)

Recommended Treatment Method:
C = Container Grown
F = Field Grown

Scientific Name	Common Name				
<i>Bougainvillea</i> spp.	barbara karst	C, F	<i>Dalea greggii</i>	trailing indigo bush	C, F
	California gold	C, F	<i>Daphne cneorum</i>	rose daphne	C, F
	pink pixie	C, F	<i>Daphne odora</i>	fragrant daphne	C, F
	scarlet o'hara	C, F	<i>Deutzia crenata</i>	nakiana-dwarf deutzia	C, F
	temple fire	C, F	<i>Deutzia gracilis</i>	slender gracilis	C, F
	Texas dawn	C, F	<i>Dodonea viscosa</i>	hopseed bush	F
<i>Buxus microphylla japonica</i>	boxwood, Japanese	F	<i>Enkianthus companulatus</i>	red-veined enkianthus	C, F
<i>Buxus microphylla Koreana</i>	Korean boxwood	F	<i>Elaeagnus pungens</i>	fruitland silver berry	C, F
<i>Buxus sempervirens</i>	boxwood, common	C, F	<i>Erica cinerea</i>	purple bell heather	C, F
<i>Buxus x 'Green velvet'</i>	green velvet boxwood	C, F	<i>Erica vagans</i>	cornish heather	C, F
<i>Callistemon citrinus</i>	bottlebrush, lemon	F	<i>Erica x darleyensa</i>	Mediterranean pink heather	C, F
<i>Callistemon viminalis</i>	weeping bottlebrush	C, F	<i>Escalonia</i> spp.	escallonia	C, F
<i>Calluna vulgaris</i>	spring torch Scotch heather	C, F	<i>Eugenia myrtifolia</i>	teenie genie brushcherry	C, F
<i>Camellia japonica</i>	camellia	C, F	<i>Eugenia myrtifolia 'Globulus'</i>	dwarf brush cherry	C, F
<i>Caryopteris clandonensis</i>	blue mist bluebeard	C, F	<i>Euonymus fortunei</i>	canadale gold euonymus	C, F
<i>Caryopteris x clandonen 'Blk night'</i>	dark knight bluebeard	C, F		emerald'n gold euonymus	F
<i>Cassia artemisioides</i>	cassia, feathery	C, F	<i>Euonymus japonica</i>	sunspot euonymus	C, F
<i>Cassia, eremophila</i>	senna	C, F		silver king euonymus	F
<i>Ceanothus</i> spp.	wild lilac	F		chollipo euonymus	C, F
<i>Cephalotaxus drupacae</i>	plum yew	C, F	<i>Euonymus kiatschovica</i>	gold spot euonymus	C, F
<i>Cerastium tomentosum</i>	snow-in-summer	C, F	<i>Euonymus vegetus</i>	silver princess euonymus	C, F
<i>Cerastostigma plumbaginoides</i>	dwarf plumbago	C, F	<i>Fatsia japonica</i>	variegated evergreen euonymus	C, F
<i>Ceratosigma willmottianum</i>	Chinese plumbago	C, F	<i>Felicia amelloides</i>	spreading euonymus	C, F
<i>Chaenomeles japonica</i>	orange flowering quince	C, F	<i>Forsythia intermedia</i>	bigleaf wintercreeper	C, F
<i>Chamaecyparis obtusa</i>	kosteri cypress	C, F	<i>Forsythia x 'Spring glory'</i>	Japanese aralia	C, F
	nana-dwarf hinoki cypress	C, F	<i>Forsythia x "Santa Claus"</i>	blue marguerite	C, F
	torulosa cypress	C, F	<i>Gardenia jasminoides</i>	forsoythia, border	C, F
<i>Chamaecyparis pisifera</i>	baileyi-dogwood	F		spring glory forsoythia	C, F
	flaviramea-dogwood	F	<i>Gaultheria procumbens</i>	santa claus fuchsia	C, F
	sawara-false cypress	F	<i>Gaultheria shallon</i>	august beauty gardenia	C, F
	squarrosa minima cypress	C, F	<i>Gelsemium sempervirens</i>	gardenia	C, F
	filifera-thread cypress	C, F	<i>Genista pilosa</i>	miniature gardenia	C, F
<i>Chrysalidocarpus lutescens</i>	areca palm	F	<i>Hamamelis virginiana</i>	radican gardenia	C, F
<i>Clethra alnifolia</i>	summersweet	C, F	<i>Hardenbergia violacea</i>	wintergreen	C, F
<i>Cleyera japonica</i>	cleyera, Japanese	C, F	<i>Hebe buxifolia</i>	salal/lemon leaf	C, F
<i>Colonema pulchrum</i>	pink breath of heaven	C, F	<i>Hibiscus rosa-sinensis</i>	carolina jessamine	C, F
<i>Convolvus cneorum</i>	bush morning glory	C, F	<i>Hibiscus syriacus</i>	woadwaxen	C, F
<i>Convolvulus mauritanicus</i>	ground morningglory	C, F		common witch hazel	C, F
<i>Cornus alba</i>	sibirica-Siberian dogwood	C, F	<i>Ilex aquifolium</i>	lilac vine	C, F
<i>Cornus stolonifera</i>	baileyi-red-osier dogwood	F		boxleaf hebe	C, F
	flaviramea-yellowtwig dogwood	F	<i>Ilex aquipernyi</i>	ross estey-hibiscus	C, F
<i>Corylus contorta</i>	walking stick	C, F	<i>Ilex attenuata</i>	rose of sharon, red bird	C, F
<i>Cotinus coggygria</i>	royal purple smoke tree	C, F	<i>Ilex cornuta</i>	rose of sharon, red heart	F
<i>Cotinus dammeri</i>	coral beauty smoke tree	C, F		rose of sharon, woodbridge	C, F
	eichholz smoke tree	C, F		Balkans holly	F
<i>Cotoneaster adpressus</i>	praecox-early cotoneaster	C, F		gold coast holly	F
<i>Cotoneaster apiculatus</i>	cotoneaster, cranberry	C, F	<i>Ilex crenata</i>	san jose holly	C, F
<i>Cotoneaster congestus</i>	cotoneaster, Pyrenees	C, F		savannah holly	C, F
<i>Cotoneaster dammeri</i>	cotoneaster, bearberry	C, F		burford holly	C, F
<i>Cotoneaster himalayan</i>	Himalayan cotoneaster	C, F		dwarf burford holly	C, F
<i>Cotoneaster horizontalis</i>	cotoneaster, rock	C, F		needle point holly	C, F
<i>Cycas revoluta</i>	sago palm	C, F		holly, Chinese	
<i>Cytisus praecox</i>	hollandia-warminster broom	C, F		compacta-dwarf Japanese holly	C, F
<i>Cytisus scoparius</i>	lena-Scotch broom	C, F		convexa holly	C, F
<i>Cytisus spp.</i>	holandia-Scotch broom	F	<i>Ilex glabra</i>	dwarf Chinese holly	C, F
	allgold warminster broom	C, F	<i>Ilex glabra</i>	green luster holly	C, F
	lilac time broom	C, F		helleri-heller's Japanese holly	C, F
				hetzii's Japanese holly	C, F
				stokesii Japanese holly	C, F
				compacta-compact inkberry holly	C, F
				nordica-inkberry holly	C, F

Shrubs (Cont.)

Recommended Treatment Method:
C = Container Grown
F = Field Grown

Scientific Name	Common Name				
<i>Ilex meserveae</i>	blue boy holly	C, F	<i>Lonicera periclymenum</i>	flowering woodbine	C, F
	blue girl holly	C, F		serotina woodbine	C, F
	China boy holly	C, F	<i>Lonicera sempervirens</i>	trumpet honeysuckle	C, F
	China girl holly	C, F	<i>Lorpetalum chinense</i>	sizzling pink fringe flower	C, F
	ebony magic holly	F	<i>Loropetalum chin</i>	razzleberri fringe flower	C, F
<i>Ilex vomitoria</i>	nana-dwarf yaupon holly	C, F	<i>rubrum 'Razzleb'</i>		
	pendula-weeping yaupon holly	C, F	<i>Mahonia aquifolium compacta</i>	dwarf Oregon grape	C, F
	yaupon holly	C, F	<i>Mahonia bealei</i>	leather leaf mahonia	C, F
<i>Illicium annisatum</i>	mystery gardenia	C, F	<i>Mahonia repens</i>	creeping mahonia	C, F
<i>Itea ilicifolia</i>	henry garnet holly leaf sweetspire	C, F	<i>Michelia figo</i>	banana shrub	C, F
<i>Ixora collinea</i>	ixora	C, F	<i>Myrica cerifera</i>	wax myrtle	C, F
<i>Juniperus chinensis</i>	hollywood juniper	C, F	<i>Myoporum parvifolium</i>	putah creek	C, F
	media-old gold juniper	C, F	<i>Nandina domestica</i>	compacta-dwarf heavenly bamboo	C, F
	pfitzer juniper	C, F		harbour dwarf-heavenly bamboo	C, F
	pfitzerana glauca-blue juniper	C, F		heavenly bamboo (nandina)	C, F
	pfitzerana-pfitzer juniper	C, F		nana compacta-heavenly bamboo	C, F
	sea green juniper	F	<i>Nerium oleander</i>	hana purpurea-heavenly bamboo	C, F
	torulosa-hollywood juniper	C, F		woods dwarf-heavenly bamboo	C, F
<i>Juniperus conferta</i>	emerald sea shore juniper	C, F		hardy red oleander	C, F
	shore juniper	C, F		oleander	C, F
<i>Juniperus horizontalis</i>	andorra juniper	C, F	<i>Osmanthus fortunei</i>	ruby lace oleander	C, F
	bar harbor juniper	C, F		fortunes osmanthus	C, F
	blue chip juniper	C, F	<i>Osmanthus fragrans</i>	sweet olive	C, F
	blue rug juniper	C, F	<i>Pennisetum setaceum 'Rubrum'</i>	purple fountain grass	C, F
	creeping juniper	C, F	<i>Phoenix roebelenii</i>	pigmy date palm	C, F
	dwarf andorra juniper	C, F	<i>Photinia fraseri</i>	graser's photinia	C, F
	huntington blue juniper	C, F	<i>Physocarpus opulifolius</i>	fwarf ninebark	C, F
	plumosa-andorra juniper	C, F	<i>Pieris japonica</i>	lily-of-the-valley	C, F
	wiltonii-blue carpet juniper	C, F		mountain fire lily-of-the-valley	C, F
<i>Juniperus procumbens</i>	nana-dwarf Japaneses	C, F		snowdrift lily-of-the-valley	C, F
	garden juniper		<i>Pieris japonica x forestii</i>	temple bells lily-of-the-valley	C, F
<i>Juniperus prostrata</i>	prostrata juniper	C, F		valley rose lily-of-the-valley	C, F
<i>Juniperus sabina</i>	broadmoor juniper	C, F	<i>Pinus mugo</i>	valley valentine lily-of-the-valley	C, F
	foemina-hicks juniper	C, F	<i>Pittosporum tobira</i>	forest flame lily-of-the-valley	C, F
	savin juniper	C, F		mugo-mugho pine	C, F
	tamariscifolia-tam juniper	C, F		green pittosporum	C, F
<i>Juniperus scopulorum</i>	emerald green juniper	F	<i>Plumbago ariculata</i>	wheeler's dwarf pittosporum	C, F
	wichita blue juniper	C, F	<i>Plumbago capensis</i>	blue cape plumbago	F
<i>Juniperus squamata</i>	blue juniper	C, F	<i>Podocarpus macrophyllus</i>	plumbago	C, F
	blue star juniper	C, F	<i>Polygala fruticosa</i>	yewpine	C, F
	parsonii juniper	C, F	<i>Polygala fruticosa</i>	sweet pea shrub	C, F
<i>Kalmia latifolia</i>	laurel, mountain	C, F	<i>Polystichum polyblepharum</i>	tassel fern	C, F
<i>Lagerstroemia indica</i>	crepe myrtle	C, F	<i>Potentilla fragiformis</i>	cinquefoil	F
<i>Leucophyllum frutescens</i>	Texas sage	C, F	<i>Potentilla fruticosa</i>	cinquefoil	C, F
<i>Leucophyllum laevigatum</i>	chihuahan sage	C, F		floppy disc cinquefoil	C, F
<i>Leucothoe axillaris</i>	leucothoe, coast	C, F	<i>Potentilla spp.</i>	gold drop pontentilla	F
<i>Leucothoe fontanesiana</i>	leucothoe, drooping	C, F		goldfinger potentilla	C, F
<i>Ligustrum japonicum</i>	privet, Japanese	C, F	<i>Potentilla verna</i>	red ace potentilla	C, F
	wax ligustrum	C, F	<i>Prunus glandulosa</i>	red ace potentilla	C, F
	yellow tip ligustrum	C, F	<i>Prunus laurocerasus</i>	sunset potentilla	C, F
<i>Ligustrum lucidum</i>	privet, glossy	C, F	<i>'Otto luykens'</i>	tangerine potentilla	C, F
<i>Ligustrum ovalifolium</i>	California privet	F	<i>Psidium cattleianum</i>	cinquefoil	C, F
<i>Ligustrum texanum</i>	howardi privet	C, F	<i>Pyracantha fortuneana</i>	spring cinquefoil	C, F
	wax leaf privet	C, F		dwarf pink flowering almond	C, F
<i>Ligustrum vicaryi</i>	privet, golden	F		otto luykens English laurel	C, F
<i>Ligustrum vulgare</i>	iodense privet	C, F		strawberry guava	C, F
<i>Livistona chinensis</i>	Chinese fountain palm	F		lolendei monrovia pyracantha	C, F
<i>Lonicera fragrantissima</i>	winter honeysuckle	C, F		monon pyracantha	F
				red elf hybrid pyracantha	F
				rutgers hybrid pyracantha	C, F
				santa cruz pyracantha	C, F
				victory pyracantha	F

Shrubs (Cont.)

Recommended Treatment Method:
C = Container Grown
F = Field Grown

Scientific Name	Common Name		<i>Rhododendron spp.</i> <i>hybrids (Cont.)</i>		
<i>Rhaphiolepis indica</i>	charisma-monruce rhaphiolepis	C, F		hot shot azalea	C, F
	enchantress-monech rhaphiolepis	F		hume azalea	F
	rhaphiolepis (India hawthorn)	C, F	<i>Rhus lancea</i>	inga azalea	F
	springtime-Monme rhaphiolepis	F	<i>Rhus typhina</i>	irene koster azalea	C, F
<i>Rhaphiolepis ovata</i>	roundleaf rhaphiolepis	C, F	<i>Rosa x 'Flower carpet'</i>	president clay azalea	C, F
<i>Rhododendron calendulaceum</i>	cannon's double azalea	C, F	<i>Rosa rugosa</i>	tradition azalea	C, F
	flame azalea	F	<i>Rosmarinus officinalis</i>	sumac, African	C, F
	golden flare azalea	C, F	<i>Senecio cineraria</i>	staghorn sumac	C, F
	Klondike azalea	C, F	<i>Skimmia japonica</i>	red groundcover rose	C, F
<i>Rhododendron campylocarpum</i>	butterfly rhododendron	F	<i>Skimmia revesiana</i>	ramanas rose	C, F
<i>Rhododendron carolinianum</i>	PJM rhododendron	C, F	<i>Solanum rantonetii</i>	rosemary	F
<i>x daurium</i>			<i>'Royal purple'</i>	dusty-miller	
<i>Rhododendron catawbiense</i>	catawba album rhododendron	C, F	<i>Spiraea bumalda</i>	Japanese skimmia	C, F
	catawba rhododendron	C, F	<i>Spiraea cinerea</i>	reeve's skimmia	C, F
	lord roberts rhododendron	C, F	<i>Spiraea japonica</i>	Paraguay nightshade	C, F
	rocket rhododendron	C, F			
<i>Rhododendron caucasicum</i>	cuninham white rhododendron	C, F		anthony waterer spiraea	C, F
<i>x ponticum</i>				first snow spiraea	C, F
<i>Rhododendron exbury</i>	cannon's double azalea	C, F	<i>Spiraea vanhouttei</i>	dolchia spiraea	C, F
	golden flare azalea	C, F	<i>Streptosolen jamesonii</i>	Japanese alpine spiraea	C, F
	Klondike azalea	C, F	<i>Syringa rothomagensis</i>	magic carpet spiraea	C, F
<i>Rhododendron forrestii repens</i>	gomer waterer rhododendron	C, F	<i>Syringa vulgaris</i>	neon flash spiraea	C, F
<i>Rhododendron forrestii</i>	elizabeth rhododendron	C, F	<i>Taxus cuspidata</i>	shirobana spiraea	C, F
<i>x griersonianum</i>			<i>Tecomaria capensis</i>	bridal wreath	C, F
<i>Rhododendron griffithianum</i>	jean marie rhododendron	C, F	<i>Ternstroemia gymnanthera</i>	marmalade bush	C, F
<i>Rhododendron hybrid spp.</i>	America rhododendron	C, F	<i>Teucrium fruticans</i>	Chinese lilac	C, F
	English Roseum rhododendron	F	<i>Thevetia nerifolia</i>	lilac, common	F
	nova zembla rhododendron	C, F	<i>Thuja occidentalis</i>	yew, Japanese	F
	scintillation rhododendron	C, F		cape honeysuckle	C, F
<i>Rhododendron impeditum</i>	rhododendron	C, F		ternstroemia, Japanese	C, F
<i>Rhododendron indica</i>	formosa azalea	C, F		bush germander	C, F
	waucabusa azalea	C, F		yellow oleander	C, F
<i>Rhododendron kaempferi</i>	blue danube azalea	C, F		emerald arborvitae	C, F
<i>Rhododendron kerume</i>	coral bells azalea	C, F		globosa-globe arborvitae	C, F
	hino crimson azalea	C, F		little giant-dwarf arborvitae	C, F
	hino pink azalea	C, F	<i>Thuja orientalis</i>	nigra-dark American arborvitae	C, F
	mildred azalea	C, F		pyramidalis arborvitae	C, F
	snow azalea	C, F		rheingold arborvitae	C, F
<i>Rhododendron maximum</i>	rhodie max (rosebay)	C, F	<i>Tibouchina urvilleana</i>	techny arborvitae	F
<i>Rhododendron mucronulatum</i>	rhododendron	F	<i>Veitchia merrilli</i>	woodwardii arborvitae	C, F
<i>Rhododendron obtusum</i>	hino-crimson azalea	C, F	<i>Viburnum bodnantense</i>	aureus nana-dwarf golden arborvitae	C, F
<i>Rhododendron ponticum</i>	chioniodes rhododendron	C, F	<i>Viburnum carlesii</i>	minima glauca-dwarf arborvitae	C, F
	daphnoides rhododendron	C, F	<i>Viburnum davidii</i>	princes flower	C, F
<i>Rhododendron racemosum</i>	dwarf scarlet wonder rhododendron	C, F	<i>Viburnum japonicum</i>	Christmas palm	F
	tribly rhododendron	C, F	<i>Viburnum judd (V. x juddii)</i>	pink dawn viburnum	C, F
	unique rhododendron	C, F	<i>Viburnum lantana</i>	Koreanspice vibunum	C, F
	vulcan rhododendron	C, F	<i>Viburnum opulus sterile</i>	david viburnum	C, F
<i>Rhododendron sassthigiatim</i>	ramapo rhododendron	C, F	<i>Viburnum plicatum tomentosum</i>	viburnum	F
<i>x carolinianum</i>			<i>Viburnum setigerum</i>	viburnum	C, F
<i>Rhododendron satuski</i>	gumpo pink azalea	C, F	<i>Viburnum tinus compactum</i>	spring bouquet viburnum	F
	higasa azalea	F	<i>Viburnum trilobum</i>	cranberry bush	C, F
	reijn azalea	C, F	<i>Viburnum trilobum compactum</i>	dwarf cranberry bush	C, F
<i>Rhododendron spp. hybrids</i>	carror azalea	C, F	<i>Viburnum x pragense</i>	viburnum	C, F
	fashion azalea	C, F	<i>Weigela florida</i>	bristol ruby weigela	C, F
	gerard christina azalea	F		java red weigela	C, F
	girard roberta azalea	C, F	<i>Xylosma congestum</i>	minuet weigela	C, F
	golden flare exbury azalea	F	<i>Yucca filamentosa</i>	xylosma	F
	helmut vogel azalea	F		yucca	C, F
	hershey red azalea	F			

Groundcovers/Perennials Recommended Treatment Method:
C = Container Grown
F = Field Grown

Scientific Name	Common Name			
<i>Achillea</i> spp.	yarrow	C, F	<i>Erigeron speciosus</i>	darkest of all fleabane
<i>Achillea filipendulina</i>	moonshine-fern/leaf yarrow	C, F	'Darkest of all'	
<i>Achillea millefolium</i>	common yarrow	C, F	<i>Euryops pectinatus</i> 'Munchkin'	dwarf euryops
<i>Achillea millefolium</i> 'Paprika'	paprika yarrow	C, F	<i>Eustoma grandiflorum</i> 'Pink'	pink lisianthus
<i>Achillea tomentosa</i>	wooly yarrow	C, F	<i>Evolvulus nuttallianus</i>	blue daze
<i>Agapanthus africanus</i>	lilly of the Nile	C, F	<i>Fatsyhedra japonica</i>	Japanese aralia
	queen anne lily of the Nile	C, F	<i>Festuca ovina glauca</i>	blue fescue
<i>Agapanthus</i> 'Peter pan'	lily of the Nile	C, F	<i>Gaillardia x grandiflora</i>	blanket flower
<i>Ammophila breviligulata</i>	beechgrass	C, F	<i>Gazania</i> spp.	gazania
<i>Aptenia cordifolia</i>	red apple aptenia	C, F	<i>Geranium cinerium</i> "Ballerina"	ballerina cranesbill
<i>Aquilegia x</i> 'Dragon fly'	columbine	C, F	<i>Geranium sanguineum</i>	bloody cranesbill
<i>Arctotheca calendula</i>	cape weed	F	'Bloody cran'	
<i>Argyranthemum frutescens</i>	butterfly argyranthemum	C, F	<i>Geranium subcaulescens</i>	black eyed magenta cranesbill
'Buterfly'			<i>Geum</i> spp	avens
<i>Asparagus densiflorus</i> 'Myers'	pony tail fern	C, F	<i>Geum quellyon</i>	geum
<i>Asparagus retrofractus</i>		C, F	<i>Gypsophila paniculata</i>	baby's breath
<i>Asparagus varieiegata</i>	tree fern	C, F	<i>Hakonechloa macroaureola</i>	golden hakonechloa
<i>Asparagus var.</i> 'Meegers'		C, F	<i>Hedera canariensis</i>	ivy, Algerian
<i>Aspidistra elatior</i>	cast iron plant	C, F	<i>Hedera helix</i>	ivy, English
<i>Aster novae-angliae</i>	New England aster	C, F	<i>Helichrysum petiolare</i>	white licorice helichrysum
<i>Aster novi-belgii</i>	New York aster	C, F	'White licorice'	
<i>Aster novi-belgii</i> 'Persian rose'	Persian rose dwarf aster	C, F	<i>Heliotropium fragrans</i>	common heliotrope
<i>Begonia sepmerflorens</i>	white ambassador begonia	C, F	<i>Hemerocallis</i> spp.	daylily
'Amb white'			<i>Hesperaloe parvifolia</i>	red yucca
<i>Bergenia cordifolia</i>	heartleaf bergenia	C, F	<i>Heuchera x</i> 'Bressingham'	bressingham coral bells
<i>Bidens ferulifolia</i> 'Peters gold'	peter's gold bidens	C, F	<i>Heuchera micrantha</i>	coral bells
<i>Brachycome x</i> 'New amethyst'	swan river daisy new amethyst	C, F	<i>Hosta</i> 'Francee'	francee plantain lily
<i>Callistepheus chinensis</i>	China aster	C, F	<i>Hosta fortunei</i>	plantain lily
<i>Carex alba</i>	frosty curls sedge	C, F	<i>Hosta lancifolia</i>	albo-marginata hosta
<i>Carex</i> spp.	variegated carex	C, F	<i>Hosta x</i> 'Patriot'	narrow leafed plantain lily
<i>Carpobrotus edulis</i>	ice plant, largeleaf	F	<i>Houttuynia cordata</i>	patriot plantain lily
<i>Catharanthus roseus</i>	Madagascar periwinkle	C, F	'Chameleon'	chameleon houttuynia
<i>Chasmanthium latifloium</i>	northern sea oats	C, F	<i>Hymenoxys acaulis</i>	angelita daisy
<i>Chrysanthemum maximum</i>	shasta daisy	C, F	<i>Hypericum</i> spp.	St. Johnswort
<i>Chrysanthemum</i> spp.	chrysanthemum species	C, F	<i>Impatiens walleryana</i> 'Lipstick'	lipstick impatiens
<i>Clivia miniata</i> 'French hybrid'	kafir lily	C, F	<i>Ipomea acuminata</i> 'Blue dawn'	blue dawn morning glory
<i>Coreopsis verticillata</i>	coreopsis, threadleaf	C, F	<i>Iris pumila</i> 'Yellow'	yellow dwarf bearded iris
<i>Coreopsis verticillata</i>	moonbeam coreopsis	C, F	<i>Iris siberica</i>	iris
'Moonbeam'			<i>Jasminum nitidum</i>	angelwing jasmine
<i>Cortaderia selloana</i>	pampas grass	C, F	<i>Jasminum polyanthum</i>	pink jasmine
<i>Cuphea hyssopifolia</i>	false or Mexican heather	C, F	<i>Lampranthus spectabilis</i>	trailing iceplant
<i>Cyperus albostratus</i>	dwarf umbrella grass	C, F	<i>Leptospermum scoparium</i>	broom teatree/manuka
<i>Dahlia hybrid Dwarf</i>	dwarf dahlia	C, F	<i>Limonium perezii</i>	statice
<i>Dahlia x</i> 'Royal Dahlietta pink'	dwarf dahlia wendy pink	C, F	<i>Liriope gigantea</i>	white lily turf
<i>Delosperma alba</i>	white iceplant	F		giant lily turf
<i>Delosperma cooperi</i>	ice plant	C, F	<i>Liriope muscari</i>	lilac beauty lily turf
<i>Delosperma nubigenum</i>	hardy ice plant	C, F		majestic lily turf
<i>Descampsia caespitosa</i>	descampsia	C, F		monroe white lily turf
<i>Dianthus gratianopolitanus</i>	crimson treasure cheddar pink	C, F	<i>Liriope spicata</i>	silvery sunproof lily turf
'Treasure'				variegated liriope lily turf
<i>Dianthus plumaris</i>	cottage pink	C, F		big blue lily turf
<i>Dietes vegeta</i>	fortnight lily	C, F	<i>Lonicera japonica</i>	green/creeping lily turf
<i>Drosanthemum floribundum</i>	trailing rosea iceplant	F	<i>Lysimachia mummularia</i>	silver dragon lily turf
<i>Drosantheumum hispidum</i>	iceplant	C, F	<i>Lysimachia punctata</i>	honeysuckle, Japanese
<i>Dymondia margaritae</i>	diamond marguerite	C, F	<i>Matthiola incana</i> 'Harmony'	moneywort
<i>Ensete ventricosum</i>	absynnian banana	C, F	<i>Miscanthus sinensis</i>	dotted loosestrife
<i>Equisetum scirpoides</i>	dwarf horsetail	C, F	<i>Miscanthus sinensis</i>	stock
<i>Erianthus ravennae</i>	hardy pampasgrass	C, F	'Gracillimus'	eulalia grass
			<i>Moraea iridioides</i>	maiden grass
			<i>Oenothera missouriensis</i>	ozark sundrops

Groundcovers/ Perennials (Cont.)

Scientific Name	Common Name	Recommended Treatment Method:	
		C = Container Grown	F = Field Grown
<i>Oenothera speciosa</i> "Siskiyou pink"	siskiyou evening primrose	C, F	
<i>Ophiopogon japonicus</i>	dwarf mondo grass	C, F	
	mndo grass	C, F	
<i>Origanum libanoticum</i>	oregano	C, F	
<i>Osteospermum fruticosum</i>	daisy, trailing African	F	
<i>Pachysandra terminalis</i>	Japanese spurge	C, F	
<i>Parthenocissus quinquefolia</i>	virginia creeper	C, F	
<i>Pelargonium x hortorum</i>	zonal geranium	C, F	
<i>Pelargonium peltatum</i>	ivy geranium	C, F	
<i>Pennisetum alopecuroides</i>	fountain grass	C, F	
<i>Pennisetum setaceum</i>	chrimson fountaingrass	C, F	
<i>Penstemon x 'Apple blossom'</i>	apple blossom penstemon	C, F	
<i>Pentas lanceolata</i>	star clusters	C, F	
<i>Perovskia atriplicifolia</i>	Russian sage	C, F	
<i>Petunia-hybrids</i>	garden petunias	C, F	
<i>Phalaris arundinacea picta</i>	ribbon grass	C, F	
<i>Phlox subulata</i>	moss pink	C, F	
<i>Polystichum polyblepharum</i>	tassel fern	C, F	
<i>Ratbida columnifera</i>	Mexican hat	C, F	
<i>Rudbeckia fulgida</i>	blackeyed susan	C, F	
<i>Ruscus hypophyllum</i>	butcher's broom (Israeli ruscus)	C, F	
<i>Salvia daghestanica</i>	daghestan sage	C, F	
<i>Salvia grahamii</i>	graham's sage	C, F	
<i>Sasa pygmaea</i>	dwarf bamboo	C, F	
<i>Scutellaria resinosa</i>	skull cap	C, F	
<i>Sedum x 'Autumn joy'</i>	autumn joy stonecrop	C, F	
<i>Sedum x 'Vera Jameson'</i>	vera jameson stonecrop	C, F	
<i>Tagetes patula 'Little Hero'</i>	little hero marigold	C, F	
<i>Trachelospermum asiaticum</i>	asaian jasmine	C, F	
<i>Tulbaghia violacea</i>	society garlic	C, F	
<i>Verbena rigida</i>	veined verbena	C, F	
<i>Vinca major</i>	periwinkle, bigleaf	C, F	
<i>Vinca minor</i>	periwinkle, dwarf	F	
<i>Vinca spp.</i>	periwinkle	F	

Non-Bearing Trees and Vines¹

	Recommended Treatment Method:	
	C = Container Grown	F = Field Grown
almond		F
apple		F
apricot		F
avocado		F
blackberry		F
blueberry		F
boysenberry		F
cherry, sour		F
cherry, sweet		F
currant		F
dewberry		F
elderberry		F
fig		F
filbert		F
gooseberry		F
grape, American		F
grape, European		F
grapefruit		F
kiwi		F

lemon	F
loganberry	F
macadamia nut	F
nectarine	F
olive	F
orange	F
peach	F
pear	F
pecan	F
pistachio	F
plum	F
pomegranate	F
prune	F
raspberry	F
walnut, black	F
walnut, English	F

¹ Non-bearing fruit and nut trees and non-bearing vineyards are defined as plants that will not bear fruit for at least one year after treatment.

Ornamental Bulbs in Landscape Settings

Gallery 75 Dry Flowable may be applied for control of susceptible annual weeds in ornamental bulbs such as bulbous iris, daffodil (narcissus), gladiolus, hyacinth, lilies, and tulip. Apply Gallery 75 Dry Flowable to the soil surface 2 to 4 weeks after planting but prior to the emergence of annual weeds. Gallery 75 Dry Flowable may also be applied following bulb emergence but prior to flowering and after flowering. For fall planted bulbs, apply Gallery 75 Dry Flowable in late winter or early spring to weed-free soil surfaces. For bulbs, make a single application within 30 days following planting and prior to bulb emergence. Do not exceed the 0.66 lb of Gallery 75 Dry Flowable per acre rate.

Use Precautions:

- Do not use Gallery 75 Dry Flowable for weed control in ornamental bulbs grown for commercial production.
- Gallery 75 Dry Flowable is not recommended for application to:
 - Tulip plants that have emerged to a height greater than 3/4 inch.
 - Gladiolus prior to emergence or if corms are less than one inch in diameter.
 - Bulbs while they are flowering.

Shadehouse Areas

Gallery 75 Dry Flowable may be applied in open shadehouse-type structures where the natural flow of air is unimpeded. Do not apply in enclosed greenhouses or in enclosed shadehouse-type structures. Do not apply within three weeks prior to enclosing greenhouses or poly-type structures.

Christmas Tree/Conifer Plantations

Gallery 75 Dry Flowable - Alone

Apply Gallery 75 Dry Flowable as a directed spray to the soil surface or as an over the top spray to established plantings of field grown Christmas tree/conifer species listed in this label. Follow all instructions provided in the General Information section of this label.

Use Precautions:

Do not apply Gallery 75 Dry Flowable to seedbeds or seedling transplant beds. Apply only to established plantings. Established plants are defined as those that have been transplanted into their final growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation.

Tank Mix Combinations

Tank mix combinations of Gallery 75 Dry Flowable plus other labeled herbicides may be used in established Christmas tree plantings. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions and limitations before use. Refer to instructions for Gallery 75 Dry Flowable in Tank Mix in the Mixing Directions section.

Gallery 75 Dry Flowable/Roundup: Apply tank mix combinations of Gallery 75 Dry Flowable plus Roundup herbicide only as directed sprays in Christmas tree plantings. When applied as directed, Gallery 75 Dry Flowable plus Roundup will provide postemergence control of susceptible weed species listed on the label for Roundup and residual preemergence control of susceptible weed species listed on the label for Gallery 75 Dry Flowable. Refer to the label for Roundup for specific use directions, precautions and limitations before use. Refer to instructions for Gallery 75 Dry Flowable in Tank Mix in the Mixing Directions section.

Use Precautions:

- Do not apply sprays containing Roundup over the top of Christmas tree plantings.
- Extreme care must be exercised to prevent contact of sprays containing Roundup with foliage or stems of Christmas trees or other desirable plants or severe plant damage or death may result.

Non-Cropland Areas

Gallery 75 Dry Flowable is recommended as a preemergence herbicide for control of listed broadleaf weeds in non-cropland areas including industrial sites, utility substations, parking lots, driveways, walking paths, highway guardrails, sign posts and delineators where maintenance of bare ground is desired.

Apply Gallery 75 Dry Flowable anytime prior to germination of target weeds. Areas to be treated should be free of established weeds or existing weeds should be controlled with postemergence herbicides.

Refer to the General Information section prior to using this product on non-cropland.

Gallery 75 Dry Flowable is compatible and can be tank mixed with other herbicides registered for use on non-cropland. Applied as directed, tank mixes containing Gallery 75 Dry Flowable will provide control of susceptible weed species listed on the respective labels. All directions, precautions and limitations on the respective product labels apply to the tank mix use. Refer to instructions for Gallery 75 Dry Flowable in Tank Mix in the Mixing Directions section.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC

Produced for Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Label Code: D02-081-019

Replaces Label: D02-081-018

LOES Number: 010-00064

EPA accepted 06/11/03

Revisions:

1. Removed lantana.