

# Roundup CUSTOM<sup>®</sup>

**FOR AQUATIC & TERRESTRIAL USE**



**A broad-spectrum postemergence herbicide for aquatic and industrial, turf, ornamental, forestry, roadside, utility rights-of-way, select crop, and other listed terrestrial weed control.**

(For a complete list of aquatic and terrestrial use sites, see the Directions for Use section of the attached labeling.)

**ACTIVE INGREDIENT:**  
 \*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt ..... 53.8%  
 OTHER INGREDIENTS: ..... 46.2%  
 100.0%

\* Contains 648 grams of the active ingredient glyphosate, in the form of its isopropylamine salt per liter, or 5.4 pounds per U.S. gallon, which is equivalent to 480 grams of the acid, glyphosate, per liter or 4.0 pounds per U.S. gallon (39.9% by weight).

EPA Reg. No. 524-343 EPA Est. 524-IA-1

**Keep Out of Reach of Children  
CAUTION**

See inside for additional Precautions.  
**COMPLETE DIRECTIONS FOR USE**

Not all products listed on this label are registered for use in California. Check the registration status of each product in California before using.

**FOR PRODUCT INFORMATION OR ASSISTANCE USING THIS PRODUCT, CALL TOLL-FREE, 1-866-99BAYER (1-866-992-2937)**

**IN CASE OF AN EMERGENCY INVOLVING THIS PRODUCT OR FOR MEDICAL ASSISTANCE, CALL COLLECT, DAY OR NIGHT, 1-800-334-7577**

Packed for:  
 BAYER CROPSCIENCE LP  
 800 N. LINDBERGH BLVD.  
 ST. LOUIS, MISSOURI 63167 USA  
 ©2020 Bayer Group. All rights reserved.

### 3.3 Physical or Chemical Hazards

Spray solution of this product may be mixed, stored and applied using stainless steel, fiberglass, plastic or plastic-lined steel containers. Do NOT MIX STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solution of this product react with such containers and tanks to produce hydrogen gas which can form a highly combustible gas mixture. This gas mixture could flash or explode if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source and cause serious personal injury.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product may only be used in accordance with the Directions for Use on this label or on separately published supplemental labeling. Supplemental labeling for this product can be obtained from your Authorized Bayer CropScience LP Retailer or Bayer CropScience LP Representative.

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard (40 CFR Part 170). This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this label only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, shoes plus socks, and chemical-resistant gloves made of any waterproof material.

### Non-Agricultural Use Requirements

The requirements in this label apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep people and pets off treated areas until spray solution has dried.

12	12.0	WEEDS CONTROLLED	11
12.1	Weed Control, Renovation and Chemical Mowing in Turf	11	
12.2	Annual Weeds	12	
12.3	Perennial Weeds	12	
12.4	Woody Brush, Trees and Vines	14	
13	13.0	LIMIT OF WARRANTY AND LIABILITY	16

### 1.0 INGREDIENTS

**ACTIVE INGREDIENT:**  
Glyphosate, N-(phosphonomethyl)glycine, in the form of its potassium salt..... 53.6%  
Surfactant..... 46.2%  
OTHER INGREDIENTS..... 100.0%

\* Contains 6.5 grams of the active ingredient glyphosate, in the form of its potassium salt, per liter or 6.5 pounds per U.S. gallon, which is equivalent to 450 grams of the acid glyphosate per liter or 410 pounds per U.S. gallon (95.9% or weight).

### 2.0 IMPORTANT PHONE NUMBERS

1. FOR PRODUCT INFORMATION OR ASSISTANCE USING THIS PRODUCT, CALL TOLL FREE, 1-866-598-6747 (1-866-592-7931)
2. IN CASE OF AN EMERGENCY INVOLVING THIS PRODUCT OR FOR MEDICAL ASSISTANCE, CALL COLLECT TOLL FREE NUMBER, 1-800-334-7577

### 3.0 PRECAUTIONARY STATEMENTS

#### 3.1 Hazards to Humans and Domestic Animals

Keep out of reach of children  
**CAUTION**

**DOMESTIC ANIMALS:** This product is considered to be relatively nontoxic to dogs and other domestic animals. However, ingestion of this product or large amounts of freshly sprayed vegetation could result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

#### Personal Protective Equipment (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, socks and shoes.

Follow manufacturer's instructions for cleaning/maintaining PPE. (Personal Protective Equipment) If there are no instructions for washables use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothes.

### 3.2 Environmental Hazards

Killing aquatic weeds can result in depletion or loss of oxygen in the water due to decomposition of dead plant material. This oxygen loss can cause fish mortalities. Consult with your State agency with primary responsibility for regulating pesticides before applying to public waters to determine if a permit is required. For terrestrial uses, do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high-water mark. Except if applying aerially over the forest canopy. Do not contaminate water when cleaning equipment or disposing of equipment wash waters and rinsate.

Read the entire label before using this product.

Use only according to label directions.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement at the end of this labeling before buying or using. If terms are not acceptable, return at once unopened.

AVOID CONTACT OF THIS HERBICIDE WITH FORAGE GREEN STEMS EXPOSED FROM WOODY ROOTS OR TRUNK OF CROPS, DESIRABLE PLANTS AND TREES, AS SEVERE PLANT INJURY OR DESTRUCTION COULD RESULT.

THIS IS AN EVIDENCE-BASED PRODUCT. BAYER CORP. DOES NOT INTEND AND HAS NOT FORESEEN IT FOR RETRIBUTION OR REPAIRING.

### CONTENTS

1	1.0	INGREDIENTS	1
2	2.0	IMPORTANT PHONE NUMBERS	1
3	3.0	PRECAUTIONARY STATEMENTS	1
3.1	Hazards to Humans and Domestic Animals	1	
3.2	Environmental Hazards	1	
3.3	Physical or Chemical Hazards	1	
4	4.0	STORAGE AND DISPOSAL	2
5	5.0	PRODUCT INFORMATION	2
6	6.0	WEED RESISTANCE MANAGEMENT	2
6.1	Weed Management Practices	2	
6.2	Management of Glyphosate-Resistant Weeds	2	
7	7.0	MIXING	2
7.1	Mixing with Water	2	
7.2	Surfactant	3	
7.3	Tank Mixtures	3	
7.4	Tank Mixing Procedure	3	
7.5	Mixing Spray Solution Concentrations	3	
7.6	Colorants and Dyes	3	
7.7	Drift Reduction Additives	3	
8	8.0	APPLICATION EQUIPMENT AND TECHNIQUES	3
8.1	Spray Nozzle Management	3	
8.2	Aerial Application Equipment	4	
8.3	Ground Application Equipment	4	
8.4	Handheld Sprayers	5	
8.5	Selective Application Equipment	5	
8.6	Injection Systems	5	
8.7	Controlled Droplet Applicator (CEA)	6	
9	9.0	AQUATIC AND TERRESTRIAL USE SITES	6
9.1	Aquatic Sites	6	
9.2	Terrestrial Sites	6	
10	10.0	ADDITIONAL SITE MANAGEMENT INFORMATION	6
10.1	Forestry, Hardwood and Christmas Tree Site Management	6	
10.2	Native and Wildlife Habitat Management	7	
10.3	Ornamental and Production Nursery Management	7	
10.4	Commercial, Residential and Recreational Area Management	7	
10.5	Pasture Management	7	
10.6	Railroad Management	7	
10.7	Rangeland Management	7	
10.8	Recreation Management	7	
10.9	Utility Management	7	
11	11.0	CROP USES	7
11.1	Tree, Vine and Shrub Crops	7	
11.1.1	Cross-Fruit Crops	7	
11.2	Annual and Perennial Crops	7	
11.2.1	Sugarcane	10	
11.3	Grass Seed and Seed Production	10	

#### 4.0 STORAGE AND DISPOSAL

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste and re-entrainment. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage and disposal.

**PESTICIDE STORAGE:** STORE ABOVE 5°F (4.5°C) TO KEEP PRODUCT FROM CRYSTALLIZING. Do not allow this product to freeze. If allowed to crystallize, contents of larger containers to not well before use. Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep container closed to prevent spills and contamination. See individual container label for additional storage conditions, if any.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in the container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable federal, state and local regulations and procedures.

**CONTAINER HANDLING AND DISPOSAL:** Non-refillable container. Do not reuse this container to hold materials other than pesticides or dinitro herbicides (rinstate). After emptying and cleaning, it may be allowable to temporarily hold residue or other pesticide-related materials in this container. Contact your state regulatory agency to determine allowable practices in your state.

Triple rinse or pressure rinse (or equivalent) this container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix tank while rinsing or collect rinsate for later use or disposal. Insert pressure rinsing nozzle into the side of the container and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip.

Once properly rinsed, some plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest collection site, contact your chemical dealer or Bayer CropScience LP at 1-866-998BAYER (1-866-992-7257).

If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the property marked container and disposing in a sanitary landfill.

#### 5.0 PRODUCT INFORMATION

**Product Description:** This product is a postemergence, systemic herbicide that, when mixed in the spray tank with a surfactant that is approved for aquatic use, may be used for both aquatic and terrestrial weed control. This product provides broad-spectrum control of many annual and perennial weeds, woody brush, trees and vines. This product does not control submerged weeds or control residual weed control in soil. It is formulated as a water-soluble liquid that, unless otherwise directed, requires dilution with water or another carrier, and the addition of a surfactant according to label directions and intended use via before application using standard and specialized pesticide application equipment.

**Mechanism of Action:** The active ingredient in this product inhibits an enzyme found only in plants and microorganisms that is essential to the formation of specific amino acids.

**Ho Soil Activity:** This product binds tightly to soil particles and does not provide residual weed control. Weeds must be emerged at the time of application to be controlled by later application of this product. Weed seeds in the soil will not be affected by this product and will continue to germinate

in the absence of other control measures, these individuals survive, produce seed and eventually become the dominant biotype in the field (weed) population selection. The best means of reducing this selection is to use diverse weed control practices such as multiple herbicides with different mechanisms of action, and often in combination with various mechanical and cultural practices.

To minimize the occurrence of herbicide-resistant biotypes, including those resistant to glyphosate, implement the following weed management practices that are practical to your situation. These management practices are applicable to reduce the spread of confirmed resistant biotypes (managing susceptible resistant biotypes) and to reduce the potential for selecting for resistance in new species (proactive resistance management).

• Use a diversified approach toward weed management focused on pre-planting weed seed production and reducing the number of weed seeds in the soil.  
• Plant crops into fields that are as weed-free as possible and then keep them as weed-free as possible.  
• Plant seed that is as weed-free as possible.  
• Scout fields and application sites routinely, before and after herbicide application.

• Use multiple herbicide mechanisms of action that are effective against the most troublesome weeds at your application site and against those with known resistance.  
• Apply herbicides at application rates listed on the label when weeds are within the size range indicated on the label.  
• Emphasize cultural practices that suppress weeds by using crop competitiveness.  
• Use mechanical and biological weed management practices, where appropriate.  
• Prevent field-to-field and within-field movement of weed seed or vegetative propagules.  
• Manage weed seed at harvest and after harvest to prevent a buildup of the weed seedbank.

**6.2 Management of Glyphosate-Resistant Biotypes**  
Appropriate testing is needed to determine if a weed is resistant to glyphosate. Call 1-866-998BAYER (1-866-992-7257) or contact your Bayer CropScience LP representative to determine if resistance in any particular weed biotype has been confirmed in your area, or visit us on the Internet at [www.weedstakeoutmanagement.com](http://www.weedstakeoutmanagement.com) or [www.weedscience.org](http://www.weedscience.org). Glyphosate-resistant weeds can be controlled or managed by applying this product in combination with residual pre-emergence herbicides and/or other post-emergence herbicides labeled for control of the targeted weed in the crop being grown on the site of application for more information, see the "WEEDS CONTROLLED" section of this label.

Since the occurrence of resistant weeds is difficult to detect prior to use, Bayer CropScience LP accepts no liability for any losses that result from the failure of this product to control resistant weeds.

**7.0 MIXING**  
Spray solutions of this product may be mixed, stored and applied using clean stainless steel, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.  
Eliminate any risk of splashing the contents of the tank back into the carrier source while mixing. Use approved anti-back-splashing devices where required by state or local regulations.  
A 50-mesh mesh screen or line strainer on the spray equipment is adequate. Clean spray parts promptly after using this product by thoroughly flushing with water.

**6.0 WEED RESISTANCE MANAGEMENT**  
Glyphosate, the active ingredient in this product, is a Group 9 herbicide based on the mechanism of action classification system of the Weed Science Society of America. Any weed population can contain plants that are naturally resistant to Group 9 herbicides. Weeds resistant to Group 9 herbicides can be effectively managed by using another herbicide from a different Group (either alone or in a mixture according to label directions), by using other cultural or mechanical methods of weed control or a combination of the two. Consult your local company representative, state cooperative extension agent, professional consultant or other qualified authority to determine appropriate actions for controlling specific resistant weeds.

**6.1 Weed Management Practices**  
Resistant populations arise when rare individual plants are uncontrollable by a normal dose of a given herbicide under normal environmental conditions.

Unattached plant stems and rootstocks beneath the soil surface will also not be affected by this product.

**Biological Degradation:** Degradation of this product is primarily a biological process carried out by soil microbes.

**Stage of Weeds:** Aquatic weeds must have foliage above the water surface in order to be controlled by this product. On terrestrial sites, annual and perennial weeds are easiest to control when they are small. See the "WEEDS CONTROLLED" section of this label for more information on the control of specific weeds.

**Cultural Considerations:** Reduced weed control could result when this product is applied to annual or perennial weeds that have been mowed, grazed or cut, and have not been allowed to re-grow prior to application. Always use the highest application rate of this product within the given range when weed growth is heavy or dense, or when weeds are growing in an undisturbed (non-cultivated) area. Reduced weed control could result when this product is applied to weeds that show signs of disease or insect damage, are covered with dust, or are surviving under poor growing conditions.

**Spray Coverage:** For enhanced results, spray coverage must be uniform and complete. Do not spray foliage to the point of runoff.  
Rainfastness: Rainfall or submersion of aquatic weeds by wave action within 4 hours of application could wash this product off of the foliage and a second application might be needed for acceptable weed control. Refer to specific use sections of this label for additional information on minimum intervals required before re-application of this product.

**Time to Symptoms:** This product moves through the plant from the point of foliage contact to and into the root system. Visible effects are a gradual wilting and yellowing of the plant that progresses to complete browning and abscission of growth and deterioration of underground plant parts. Effects are visible on most annual weeds within 2 to 4 days, but on most perennial weeds, effects might not be visible for 7 or more days after application. Extremely cool or cloudy weather following application could slow activity of this product and delay development of visual symptoms.

**Maximum Application Rates:** The maximum application or use rates stated throughout this label are given in units of volume (fluid ounces or quarts) of this product per acre. However, the maximum allowable application rates apply to this product combined with the use of any and all other herbicides containing the active ingredient glyphosate, whether applied separately or in a tank mixture, on a basis of total pounds of glyphosate (total equivalent) per acre. If more than one glyphosate-containing product is applied to the same site within the same year, you must ensure that the total use of glyphosate (pounds and equivalent) does not exceed the maximum allowed. See the "INGREDIENTS" section of this label for necessary product information.

Unless otherwise specified on this label, the combined total of all applications of this product on a site must not exceed 6 quarts (6 pounds) of glyphosate and/or per acre per year.

**NOTE:** Use of this product in any manner not consistent with this label could result in injury to persons, animals, crops or other desirable vegetation, or have other unintended consequences.

**6.0 WEED RESISTANCE MANAGEMENT**  
Glyphosate, the active ingredient in this product, is a Group 9 herbicide based on the mechanism of action classification system of the Weed Science Society of America. Any weed population can contain plants that are naturally resistant to Group 9 herbicides. Weeds resistant to Group 9 herbicides can be effectively managed by using another herbicide from a different Group (either alone or in a mixture according to label directions), by using other cultural or mechanical methods of weed control or a combination of the two. Consult your local company representative, state cooperative extension agent, professional consultant or other qualified authority to determine appropriate actions for controlling specific resistant weeds.

**6.1 Weed Management Practices**  
Resistant populations arise when rare individual plants are uncontrollable by a normal dose of a given herbicide under normal environmental conditions.



responsible for considering all these factors when making decisions regarding the application of this product.

The likelihood of injury occurring as the result of spray drift while applying this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or generation of fine particles (mist) that are likely to drift.

**TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFERS MUST BE MAINTAINED**  
**AVOID APPLYING THIS PRODUCT AT EXCESSIVE SPEED OR SPRAYER PRESSURE**

## 8.2 Aerial Application Equipment

Unless otherwise prohibited, all broadcast applications of this product described on this label may be made using aerial application equipment where appropriate, provided that the applicator complies with the restrictions and restrictions specified on this label and on separate supplemental labeling published for this product.

**DO NOT APPLY THIS PRODUCT USING AERIAL APPLICATION EQUIPMENT EXCEPT UNDER CONDITIONS SPECIFIED ON THIS LABEL OR ON SEPARATELY PUBLISHED SUPPLEMENTAL LABELING FOR THIS PRODUCT**  
**FOR SPECIFIC USE, INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS RELATED TO THE AERIAL APPLICATION OF THIS PRODUCT IN CALIFORNIA OR SPECIFIC COUNTIES THEREIN, REFER TO THE LIMITATIONS ON AERIAL APPLICATION IN THAT STATE OR COUNTY PRESENTED IN THIS SECTION**

Apply this product at the rate specified on this label in 3 to 25 gallons of water per acre, unless otherwise directed. Use a larger spray volume within this range where weeds, brush, trees and vines are dense or form multiple canopy layers.

Avoid direct application to any body of water  
Drift control reduction additives may be used  
Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

### Aircraft Maintenance

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills.  
**PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES COULD RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART**  
**LANDING GEAR IS MOST SUSCEPTIBLE.** Maintaining an organic coating (paint) that meets aerospace specification MIL-C-18413 can help prevent corrosion.

### AERIAL SPRAY DRIFT MANAGEMENT

The following drift management requirements must be followed to minimize off-target drift movement during aerial application. These requirements do not apply to forestry applications.

1. The distance of the downmost nozzles on the boom must not exceed 1/3 the length of the wingspan or rotor.
2. Nozzles must always point backwards parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they must be followed.

### Importance of Droplet Size

The most effective way to reduce drift potentials is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if the application is made improperly or under unfavorable environmental conditions, such as in windy, high temperature with low humidity, and/or inversion conditions as described below.

### Controlling Droplet Size

- Volume: Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- Pressure: Operate at a spray pressure towards the lower end of the range listed for the nozzle. Higher pressure reduces droplet size and does not

improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

- **Number of nozzles:** Use the minimum number of nozzles that provides uniform coverage.

- **Nozzle orientation:** Directing nozzles so that the spray is released backwards, parallel to the air stream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

- **Nozzle type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

- **Boom length:** For some use patterns, reducing the effective boom length to less than 1/2 of the wingspan or rotor length can further reduce drift without reducing swath width.

- **Application height:** Application must be made at a height of 10 feet or less above the top of the tallest plants, unless a greater height is required for aircraft safety. Making the application at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

### Swath Adjustment

When an application is made in the presence of a crosswind, the swath will be displaced downwind. Therefore, on the upwind and downwind edges of the field, the applicator must compensate for the displacement by adjusting the path of the aircraft upwind. Increase the swath adjustment distance with increasing drift potential (in high wind, smaller droplets, etc.)

### Wind

Drift potential is lowest at wind speeds of between 2 and 10 miles per hour. However, many factors, including droplet size and equipment type, determine drift potential at any given wind speed. Avoid application when wind speeds are below 2 miles per hour due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect drift.

### Temperature and Humidity

When making an application in low relative humidity, set application equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

### Temperature Inversions

Do not apply this product during a temperature inversion as drift potential is high under these conditions. Temperature inversions restrict vertical air mixing, which causes small droplets to remain suspended in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are cleared or lifted by increasing temperatures with altitude and are common on nights with layered cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog. However, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

### Sensitive Areas

Apply this product only when the potential for drift to adjacent sensitive non-target areas (e.g. residential areas, known habitat for threatened or endangered species, crop, target crops) is minimal (e.g. when wind is blowing away from a sensitive area).

### State Specific Limitations on Aerial Application

#### LIMITATIONS ON AERIAL APPLICATION IN CALIFORNIA ONLY

**DO NOT apply this product using aerial application equipment in residential areas AVOID DRIFT - DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT. DRIFT OF THIS PRODUCT MAY CAUSE VEGETATION TO WHICH APPLICATION WAS NOT INTENDED CAN CAUSE DAMAGE TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, USE PROPER AERIAL APPLICATION EQUIPMENT FITTED WITH APPROPRIATE NOZZLES AND MAINTAIN ADEQUATE BUFFERS**

Follow the directions below when making an aerial application near non-target crops, desirable annual vegetation, or desirable perennial vegetation after leaf break and before total leaf drop.

1. Do not apply this product within 100 feet of all desirable vegetation or non-target crops.
2. If winds are blowing up to 5 miles per hour, TOWARD desirable vegetation or non-target crops, do not apply this product within 500 feet of the desirable vegetation or crops.
3. If winds are blowing between 5 and 10 miles per hour, TOWARD desirable vegetation or non-target crops, a buffer zone greater than 500 feet might be needed to protect the desirable vegetation or crops.
4. Do not apply this product using aerial application equipment when winds are blowing in excess of 10 miles per hour.
5. Do not apply this product using aerial application equipment when inversion conditions exist.

When tank mixing this product with 7.4-D only 7.4-D alone formulations may be applied in California using aerial application equipment. Tank mixtures of this product with 7.4-D alone formulations may be applied by air in California on lawns, fields, and in restored tillage systems and for pasture renovation applications only.

This product, when tank-mixed with dicamba, may not be applied by air in California.

#### ADDITIONAL LIMITATIONS ON AERIAL APPLICATION IN FRESNO COUNTY, CALIFORNIA ONLY

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

The following information applies only from February 15 through March 31 within the following boundaries of Fresno County, California:

- North Fresno County line
- South Fresno County line
- East State Highway 93
- West Fresno County line

Observe the following directions to minimize off-site movement during aerial application of this product: Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor and aerial applicator.

#### Written Directions

Written directions MUST be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. These written directions MUST state the proximity of surrounding crops and that conditions of each manufacturer's product label and that label have been satisfied.

#### Aerial Applicator Training and Equipment

Aerial application of this product is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticides Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, certified in flight and certified at a Fresno County Agricultural Commissioner approved fly-in test and carburetor spray equipment at intervals sufficient to ensure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such recalibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved fly-in test constitutes such documentation or other written records showing calculations and measurement of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

**Applications at Night** - Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.  
For additional information on the proper aerial application of this product in Fresno County, call (800) 332-3111.

## 8.3 Ground Application Equipment

Apply this product at the application rate as specified on this label in 3 to 40 gallons of water per acre when making a broadcast application using ground application equipment, unless otherwise directed on this label or on

separate supplemental labeling or fact sheets published for this product. As the weed density increases, increase the spray volume toward the upper end of this range to ensure complete coverage. Use nozzles that will avoid generating a fine mist. For enhanced results with ground application equipment, use fan-tan nozzles. Check spray pattern for uniform distribution of spray droplets.

## 8.4 Handheld Sprayers

When using a handheld sprayer, apply spray solutions of this product uniformly and completely to the foliage of target weeds using a coarse droplet spectrum and a spray-to-wet technique. Do not spray to the point of runoff. For the appropriate concentration of this product in the spray solution and for the "WEEDS CONTROLLED" section of this label, refer to the "WEEDS CONTROLLED" section of this label.

For control of annual weeds, make application when weeds are small and prior to seedhead or bud formation. For control of perennial weeds, woody brush, trees and vines, make application after flowering and before fall color and leaf drop.

When making a low-volume directed spray application to annual and perennial weeds, woody brush, trees and vines using a handheld sprayer, ensure that at least 50 to 75 percent of the foliage or the top one-half of each unwanted plant is sprayed. If a straight stream nozzle is used, start the application at the top of the targeted plant and spray from top to bottom in a lateral zig-zag motion. To ensure uniform and complete coverage, spray both sides of large or tall woody brush, trees and vines, or when foliage is thick and dense, or when there are multiple sprouts. For enhanced results on woody brush, trees and vines, apply to actively growing vegetation after full leaf expansion and flowering, prior to fall color and leaf drop.

The following table summarizes various methods of label application using a backpack sprayer with a spray-to-wet or low-volume directed spray technique and high-volume sprayer application using handheld application equipment for control or partial control of herbaceous weeds, woody brush, trees and vines listed in the "WEEDS CONTROLLED" section of this label.

Method of Application	Spray Solution Concentration	Spray Volume
Handgun or Backpack Sprayer	0.5 to 1.5% by volume	Spray-to-wet technique
Low Volume Directed Spray (Backpack)	4 to 6% by volume	15 to 25 gallons/acre
High Volume Directed Spray	1.5 to 3% by volume	40 to 60 gallons/acre

Low-volume directed spray application with a backpack sprayer works best when applying to weeds and brush less than 10 feet tall. For taller weeds and brush, a high-volume handgun can be modified by replacing the nozzle size and spray pressure to produce a modified high-volume directed spray application.

## 8.5 Selective Application Equipment

Selective application equipment allows this product to be applied to weeds growing near a crop or other desirable vegetation without killing the desirable vegetation. Selective application equipment must be capable of preventing all contact of the herbicide solution with the desirable vegetation and species without spray mist escape, leakage or dripping of the herbicide solution. AVOID CONTACT OF THIS HERBICIDE WITH DESIRABLE VEGETATION. Contact of this product with desirable vegetation could result in unwanted plant damage or destruction. To the extent consistent with applicable law, such damage shall be the sole responsibility of the applicator.

This product may be diluted with water and applied using a rescabbling sprayer, shielded sprayer, hooded sprayer, wiper applicator or sponge bar to weeds listed on this label growing in any aquatic or on any terrestrial non-crop or non-field crop site listed on this label, where feasible. This product may also be used with sprayers equipped with optical weed sensor technology

Other selective equipment that may be used to deliver or apply this product are single and twin boom injectors, tree injectors, wiper applicators for cut stem and cut stump applications, and sprayer or sprayer bottles for cut stem cut stump and fall applications to control large stem weeds, brush, trees and vines listed on this label.

### Rescabbling Sprayer

A rescabbling sprayer directs the spray solution onto weeds growing above desirable vegetation, while spray solution that is not intercepted by weeds is collected and returned to the spray tank for reapplication. A rescabbling sprayer may be used to apply spray solutions of this product to weeds listed on this label in any aquatic or on any terrestrial non-crop site described on this label.

### Shielded and Hooded Sprayers

A shielded sprayer directs the herbicide solution to the target weeds while protecting desirable vegetation from contact with the herbicide spray with an impervious material or shield. Use nozzles that provide uniform coverage within the application area. Keep shields properly adjusted to protect desirable vegetation.

A hooded sprayer is a type of shielded sprayer where the spray pattern is fully enclosed, including the top, sides, front and back, thereby shielding desirable vegetation from the spray solution.

This product may be diluted with water and, unless otherwise directed, mixed with a surfactant and applied using a shielded or hooded sprayer to weeds listed on this label growing in any aquatic or on any terrestrial non-crop site listed on this label, where feasible, and between rows of plants (row middles) in any cropping system listed on this label.

Properly adjust the hood to protect desirable vegetation. Ensure that the hood is capable of completely enclosing the spray pattern. If necessary when applying around crops grown on raised beds, extend the front and rear flaps of the hooded sprayer downward to reach the ground in deep furrows.

A hooded sprayer must be configured and operated in a manner that minimizes bouncing and weeds raising the hood up off the ground surface at any time. If the hood is raised, spray nozzles can escape and come into contact with the crop or other desirable vegetation, causing damage to or destruction of the desirable vegetation. Avoid operating this equipment on rough or sloping terrain where the spray hood is likely to rise up off the ground surface.

Use hoods designed to minimize excessive dripping or runoff down the inside of the hood, such as a single, low pressure, low drift, flat fan nozzle with an 80- to 90-degree spray angle positioned at the top center of the hood, with a spray volume of 70 to 30 gallons per acre.

The following procedures will help reduce the potential for injury to desirable vegetation when using a hooded sprayer.

- Operate the sprayer with the hood on the ground or skimming across the ground surface.
- Leave at least an 8-inch untreated strip over the drill row. (For example, if a crop row width is 38 inches, use a sprayer hood with a maximum width of 30 inches.)
- Operate at a ground speed no greater than 5 miles per hour to minimize bouncing of the hooded sprayer.
- Apply when wind speed is 10 miles per hour or less.
- Use low-drift nozzles that provide uniform coverage within the application area.

Injury to a crop or other desirable vegetation can occur when application is made to leakage of weeds that come into direct contact with the desirable vegetation. Do not apply this product when leaves of desirable vegetation are growing in direct contact with weeds. Droplets must form or splatter of the herbicide solution, setting onto desirable vegetation can result in discoloration, stunting or destruction.

### Wiper Applicator

A wiper applicator is a device that physically wets the product to solutions of this product directly onto the weed or cut stump. Any non-wetted device that is capable of physically weting this product or solutions of this product directly onto the target weed or stump, such as a paint brush, may be used

A mechanical wiper applicator, such as a rope wick or sponge bar that can be driven through a field over the top of a crop or other desirable vegetation to control weeds that are taller than the desirable vegetation, must be designed, manufactured and operated to prevent the herbicide solution from coming into contact with desirable vegetation.

Wiper applicators may be used over the top of food or feed crops. ONLY if specifically permitted for use over that crop by this label or by separately published supplemental labeling for this product.

When using a mechanical wiper applicator, adjust the height of the applicator to ensure adequate contact with the weeds and so that the wiper contact point is at least 2 inches above the crop or desirable vegetation. Enhanced results can be obtained when more of the weed is exposed to the herbicide solution and weeds are a maximum of 6 inches above the desirable vegetation. Weeds that do not come into contact with the herbicide solution will not be affected. No contact can occur when weeds are growing in dense clumps, when operating in areas of severe weed infestation or when weed height varies dramatically. In these situations, more than one application of this product might be necessary.

Operate wiper applicators at a ground speed of no greater than 5 miles per hour. Performance in areas of heavy weed infestation can be improved by reducing speed, which will provide more time for re-saturation of the wiper with the herbicide solution and more contact time of the wiper with the weed. Enhanced results with a wiper applicator can be obtained when two applications are made traveling in opposite directions in the field.

Keep wiper surfaces clean.

Droplets must form or splatter of the herbicide solution, setting onto desirable vegetation can result in discoloration, stunting or destruction. Avoid leakage or dripping onto desirable vegetation. Be aware that on sloping ground the herbicide solution can migrate to one side, causing dripping on the lower end and drying of the wiper on the upper end of the applicator.

Do not apply this product using a wiper applicator when weeds are wet.

Add a nonionic surfactant to a concentration of 10 percent by volume of the total applicator solution (one gallon of surfactant for every 10 gallons of solution) for use in a wiper applicator. See the "MIXING" section of this label for more information on the use of surfactants.

For Rope Wick and Sponge Bar Applicators, apply solutions ranging from 33 to 75 percent of this product by volume in water.

For Panel Applications, apply solutions ranging from 33 to 90 percent of this product by volume in water.

As only the amount of this product that will be used during a 1-day period, as reduced product performance can result from the use of solutions held in storage.

Clean wiper parts promptly after using this product by thoroughly flushing with water.

### Single and Hollow Stem Injectors

Control of certain weeds listed in the "WEEDS CONTROLLED" section can be obtained by injecting this concentrated product or solutions of this product directly in or onto the target weed. Ensure that the handheld injector being used for this application is capable of accurately delivering the volume specified on the label. When making stem injections, the combined total use of this product must not exceed 6 quarts per acre per year. At 5 millibars of concentrated (undiluted) product per stem, 8 quarts will treat approximately 1500 stems per acre per year. The number of stems that can be treated per acre will vary depending on the injection volume and the concentration of this product in the application solution.

## 8.6 Injection Systems

This product may be used in aerial and ground injection spray systems as a liquid concentrate or diluted prior to injecting into the spray stream. Do not use this concentrated product with the undiluted concentrate of other products when using injection systems, unless otherwise directed. A non-ionic surfactant concentration of 0.5% or more in the spray stream is required for use of this product in injection systems.



techniques that prevent or minimize contact of this product with foliage of desired trees or other plants through direct contact or off-target spray movement.

#### Conifer Release - Broadcast Application

This product may be directly applied over the top of conifer tree species listed in this section after formation of final conifer resting buds in the fall or prior to initial bud swelling in the spring for control, partial control or suppression of herbaceous weeds and hardwoods listed in the "WEEDS CONTROLLED" section of this label to facilitate the release of these tree species in a forestry plantation or nursery setting. Unless otherwise specified, make this application only where conifers have been established for a minimum of one growing season.

**PRECAUTIONS:** Conifer injury can occur when this product is applied at rates higher than prescribed on this label where spray applications overlap. If application is made when conifers are actively growing or when they are growing under stress from drought, flood, insect/pest/planting or insect, animal or disease damage.

#### Conifer Release - Broadcast Application

For release of the following conifer species growing for a minimum of one growing season in most areas within the Southeastern United States, apply 24 to 48 fluid ounces of this product per acre as a broadcast application over the top of the conifer trees.

- Douglas fir
  - Pines\*
  - Fir species
  - California redwood
  - Spruce
  - Hemlock
- \*Includes all species except loblolly pine, longleaf pine, shortleaf pine or slash pine.

Apply 24 to 40 fluid ounces of this product for release of Douglas fir, pine and spruce that have been established for only one growing season (except in California).

For release of spruce (*Picea spp.*) in Maine, Michigan, Minnesota, New Hampshire and Wisconsin, up to 25 quarts of this product may be applied after formation of final resting buds in the fall for control of woody brush and tree species.

**PRECAUTIONS:** Ensure that the conifers are well hardened off before application of this product. Some nonionic surfactants can cause tree injury when broadly applied over the top of hemlock and California redwood and in mixed conifer stands. Test the nonionic surfactant to be used for tree safety before using.

#### Conifer Release in the Southeastern United States

For release of the following conifer species established for more than one growing season in the southeastern United States, apply 36 to 60 fluid ounces of this product per acre in the fall as a broadcast application over the top of the trees. For release of these species after only one growing season, apply only 24 fluid ounces of this product per acre.

- Eastern white pine
- Shortleaf pine
- Loblolly pine
- Slash pine
- Virginia pine
- Longleaf pine

**TANK MIXTURES:** This product may be applied for conifer release in a tank mix with the following products to provide a broader spectrum of postmergence weed control and for residual control of weeds below the level of those products. Only apply these tank mixtures over the top of conifer species that are approved for this use for all products in the mix. Refer to the individual product labels for approved uses and application rates. Read and follow all directions for use and precautions for each product used. Use this product according to the most restrictive precautionary statements of any product in the mixture.

Arsenal, Arsenal Herbicide, Agalators, Concentrate, Dust Extra, Oust XP, alazone, imazapic, metolachlor, metolachlor, sulfentrazone, metolachlor.

For release of Douglas fir established for a minimum of one growing season prior to bud swell in early spring, apply 24 fluid ounces of this product in a tank-mix with an appropriate rate of alazine. Do not use surfactant for this application.

For herbaceous release of loblolly pine, Virginia pine and longleaf pine in the spring and early summer, apply 12 to 18 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Oust Extra or Oust XP.

#### Late-Summer and Fall White Pine Residue Bad Formation

For release of jack pine, white pine and white spruce, apply 24 to 48 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Oust Extra or Oust XP that will not treat these conifer species.

For release of Douglas fir, apply 24 to 36 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Arsenal or Arsenal Herbicide Agalators Concentrate.

For release of balsam fir and red spruce, apply 48 fluid ounces of this product per acre in a tank-mix with an appropriate rate of Arsenal or Arsenal Herbicide Agalators Concentrate.

### 10.2 Native and Wildlife Habitat Management

This product may be used to control exotic and other undesirable vegetation in wildlife habitat and natural areas, including riparian and estuarine areas, riparian and wildlife refuges. Application may be made to allow recovery of native plant species or prior to planting desirable native species, and for similar broad-spectrum vegetation control. Spot treatment, cut stump, cut stem, stem injection, wiper application and all other methods of application listed on this label may be used to selectively remove unwanted plants for habitat management and enhancement.

This product may also be used to eliminate annual and perennial weeds prior to planting wildlife food plots. Key wildlife food species may be placed after applying this product to native species may be allowed to regenerate the area naturally. If usage is needed to manage a seedbed, wait a minimum of 7 days after application before tilling to allow establishment of this product into underground plant parts.

### 10.3 Ornamental and Production Nursery

#### Management

All uses of this product described on this label may be used in a plant nursery setting using any method of application described.

This product may be used to clear an area of unwanted vegetation prior to placing any ornamental plant, tree, shrub or other plants.

This product may also be used to control weeds growing among established woody ornamental species, such as arborescens, acacia, birch, crabapple, eucalyptus, eucalyptus, forsythia, forsythia, holly, hick, magnolia, maple, oak, redbud, pine, spruce and yew. This product may also be used to trim and edge around potted plants and other objects in a plant nursery.

**PRECAUTIONS:** Protect desirable plants from the spray solution using sheets or coverings made of water-proof material. Take care to avoid contact of spray with or mist with foliage, green stems or ornamental bank of established ornamental species.

#### Greenhouse/Stadthouse

This product may be used to control weeds growing in and around greenhouses and stadhouses.

**PRECAUTIONS:** Desirable vegetation must not be present during application in a greenhouse. Turn air circulation fans off before applying this product inside a greenhouse or stadhouse and leave them off until the application solution has dried.

### 10.4 Commercial, Residential and Recreational Area Management

All applications of this product described on this label may be used on commercial, residential and recreational areas, including parks, schools and athletic fields using any method of application described on this label including spot treatment of unwanted vegetation, trim-and-edge application around trees, fences, walking paths, buildings, sidewalks, nature trails and other objects in these areas, to eliminate unwanted weeds growing in established shrub and ornamental beds, for best management and removal and to eliminate vegetation from a site post-development, including post-

to planting an area to ornamentals, lawns or turfgrass (seed or seed), or beginning construction projects.

### 10.5 Pasture Management

The use of this product in pastures includes use on bahiagrass, bermudagrass, bluegrass, brome, fescue, guineagrass, harrington, orchardgrass, pangola grass, ryegrass, timothy, and wheatgrass.

#### Preplant, Preemergence, Pasture Renovation

This product may be applied prior to planting or emergence of forage or perennial grasses. Refer to the "WEEDS CONTROLLED" section of this label for application rates of this product for control of specific weeds.

**RESTRICTIONS:** If the total application rate of this product is 2.25 quarts per acre or less, no waiting period between application and seeding or broadcast grazing is required. If the rate is greater than 2.25 quarts per acre, remove domestic livestock before application and wait a minimum of 6 weeks after application before grazing or harvesting.

#### Spot Treatment, Wiper Application

This product may be applied in pastures as a spot treatment or over the top of desirable grasses using a wiper applicator to control taller growing weeds. For enhanced weed control, remove domestic livestock before application to allow for sufficient plant growth and wait a minimum of 7 days after application before grazing livestock or harvesting for feed. See additional instructions on the use of wiper applicators in the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label.

**RESTRICTIONS:** For spot treatment or use with a wiper applicator at rates of 2.25 quarts per acre or less, this product may be applied over the entire pasture or any portion of it. At rates greater than 2.25 quarts per acre, this product may be applied over no more than 10 percent of the total pasture at any one time. Application may be repeated in the same area at 30-day intervals.

#### Weed Suppression in Dominant Pastures

This product may be applied in dominant pastures to suppress competitive growth and seed production of annual weeds and other undesirable vegetation. Apply 9 to 12 fluid ounces of this product per acre using broadcast application equipment on pastures in late fall after desirable perennial grasses have reached dormancy or in late winter before desirable perennial grasses break dormancy and initiate green growth.

**PRECAUTIONS:** Higher application rates may be used for hard-to-control weeds; however, higher rates can cause stand reduction. Some shading of perennial grasses can occur if broadcast application is made when they are not dormant.

**RESTRICTIONS:** No waiting period is required between application and grazing or harvesting for feed. Do not apply more than 2.25 quarts of this product per acre per year onto pasture grasses except for renovation. If re-seeding is needed due to severe stand reduction, no waiting period is required after application of this product before seeding the pasture grasses listed at the beginning of this section. For all other pasture grasses, wait a minimum of 30 days after application before seeding.

### 10.6 Railroad Management

All uses of this product described in the "WEEDS CONTROLLED" or any other section of this label may be used on railroad sites using any method of application described.

This product requires a nonionic surfactant that is labeled for the intended use on the site of application to be used on the spray mixture. If application is to be made where aquatic sites might be directly sprayed or indirectly oversprayed, the surfactant must be labeled for aquatic use. Use of this product without a surfactant will result in reduced performance. See the "TANK MIXTURES" section of this label for more information on the use of surfactants with this product.

Application of this product along railroad rights-of-way may be made in up to 60 gallons of spray solution per acre.













**PERENNIAL WEEDS RATE TABLE**

Weed Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
Bromegrass, smooth	1.5 - 2.3	0.75

Apply this product when most target plants have reached the boot to head stage of development. When application is made prior to the boot stage, reduced control can result. In the fall, make application before plants have turned brown.

<b>Barnyard, woolly leaf</b>	1.5
<b>Cynodgrass, seed</b>	1.5 - 2.3

Apply this product when most target plants have reached the boot to head stage of development. When application is made prior to the boot stage, reduced control can result. In the fall, make application before plants have turned brown.

<b>Critical</b>	2.3 - 3.75	0.75
-----------------	------------	------

Apply this product when target plants are actively growing and are at or beyond the early to full bloom stage of development. Enhanced results are achieved when application is made during the summer or fall months.

<b>Clower, red, white</b>	2.3 - 3.75
<b>Cogongrass</b>	2.3 - 3.75

Apply this product in late-summer or fall when cogongrass is at least 18 inches tall and actively growing. Due to uneven stages of growth and the dense nature of cogongrass vegetation, more than one application might be necessary to achieve control.

**Cogongrass**  
2 - 8

Prior to application of this product for control of cogongrass, survey the area to determine if shellfish beds exist within the application area. If shellfish are intended to be harvested in the area, delay application of this product until after harvest or maintain a 50-foot buffer between the application area and commercial shellfish beds, or don't harvest shellfish for a minimum of 30 days following application of this product. See restrictions below.

Local conditions for control of cogongrass are when target plants are free of soil and debris and actively growing, and good spray coverage is achievable. The presence of debris or silt on the surface of cogongrass will reduce the performance of this product. To improve herbicide uptake, wash targeted plants prior to application and allow a minimum of 4 hours for plants to dry before applying this product. Where cogongrass has been cut or mowed prior to application, allow for sufficient re-growth before applying this product to ensure adequate interception and uptake of this product. Rainfall or immersion of the plant in freshwater within 4 hours after application could reduce the effectiveness of this product.

Apply 2 to 8 quarts of this product per acre using ground broadcast application or aerial spray equipment in 5 to 100 gallons of spray solution or in 5 to 10 gallons of spray solution per acre when using aerial application equipment.

Apply a 5- to 8-percent solution of this product when using a handheld backpack sprayer or high volume sprayer. Make all applications of this product for the control of cogongrass in a spray solution containing 0.25% or more (1 or more quarts per 100 gallons of spray solution) of a nonionic surfactant or other adjuvant that is compatible with this product and labeled for use with herbicides and approved for use on aquatic sites. For enhanced results, ensure complete coverage of cogongrass clumps.

**RESTRICTIONS:** If a minimum 50-foot buffer is maintained between the application area and commercial shellfish beds, there is no restriction on shellfish harvest. If application is made within 50 feet of commercial shellfish beds, DO NOT harvest shellfish for a minimum of 14 days following application of this product.

**PERENNIAL WEEDS RATE TABLE**

Weed Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
<b>Catgrass, giant*</b>	3	1

More than one application of this product will be required to achieve control especially where vegetation is partially submerged in water. Allow target weeds to re-grow to the 7- to 10-leaf stage before making next application.

<b>Dallisgrass</b>	2.3 - 3.75	1.5
<b>Dandelion</b>	2.3 - 3.75	1.5
<b>Dock, curly</b>	2.3 - 3.75	1.5
<b>Dogbane, bump</b>	3	1.5

Apply this product when most target plants have reached the late boot to flower stage of growth. For enhanced results, make application in late-summer or fall.

<b>Fescue (except tall)</b>	2.3 - 3.75	1.5
<b>Fescue, tall</b>	2.3	1

Apply this product when most target plants have reached the boot to head stage of growth. If applied prior to the boot stage, less than desirable control might be obtained.

<b>Guinea grass</b>	2.3	0.75
---------------------	-----	------

Apply this product when most target plants have at least reached the 7 leaf growth stage.

<b>Hemlock, poison</b>	1.5 - 3	0.75 - 1.5
------------------------	---------	------------

Control can also be achieved by injecting 5 milliliters of a 5-percent solution of this product using a handheld injection device in one leaf cane per plant 12 inches above the root crown. No surfactant is required.

<b>Hoagwood, giant</b>	-	-
<b>Horsetail</b>	2.3 - 3.75	1.5
<b>Horseneach</b>	3	1.5

Apply this product when most target plants have reached the late boot to flower stage of development. For enhanced results, apply in late-summer or fall.

<b>Horsetail, field</b>	-	-
<b>Iceplant</b>	1.5	1.5

Inject 0.5 milliliters of this product per stem directly into the plant stem, one segment above the root crown. No surfactant is required.

<b>Iris, yellow flag</b>	-	-
<b>Juncus, German</b>	1.5 - 3	0.75 - 1.5
<b>Juncus, artichoke</b>	2.3 - 3.75	1.5
<b>Johannesgrass</b>	1.5 - 2.3	0.75

Apply this product when most target plants have reached the boot to head stage of development or before plants have turned brown in the fall. When applied prior to the boot stage, reduced control can result.

<b>Kilbuck grass</b>	1.5 - 2.3	0.75
----------------------	-----------	------

**PERENNIAL WEEDS RATE TABLE**

Weed Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
<b>Knockweed, Bohemian, giant</b>	3	2

Apply this product when most target plants have reached the late boot to flower stage of growth. For enhanced results, apply in late-summer or fall.

<b>Japanese Knotweed</b>	3
--------------------------	---

Apply 3 quarts of this product per acre as a broadcast application in 3 to 40 gallons of spray solution with 0.5 to 1 percent by volume of a nonionic surfactant. For application using a backpack sprayer and a spray-to-wet technique, apply a 2-percent solution of this product containing 0.5 to 2 percent by volume of a nonionic surfactant. For enhanced control, do not disturb vegetation in the application area for a minimum of 7 days after application.

Control can also be achieved by cutting stems cleanly just below the 2nd or 3rd node above the ground and immediately apply 0.35 fluid ounce (10 milliliters) of a 50-percent solution of this product in water into the "wax" or remaining internode. Focus on the upper plant material that was removed from separating nodes. Use of a bio-barrier, such as carborane, physical or chemical and property associated to prevent new plants from propagating is planned and will help guard against the spread of plant material. The combined total application rate of this product must not exceed 8 quarts per acre. Control can also be achieved by injecting 5 milliliters of this product per plant into the second or third internode using a handheld injection device. No surfactant is required.

<b>Lantern</b>	-	0.75 - 1
----------------	---	----------

Apply this product when most target plants are at or beyond the bloom stage of growth. Use the higher spray solution concentration on plants that have reached the woody stage of growth.

<b>Lespedeza</b>	2.3 - 3.75	1.5
<b>Loosestrife, purple</b>	2	1 - 1.5

Apply this product when most target plants are at or beyond the bloom stage of growth. Enhanced results can be achieved when application is made during summer or fall months. Fall application must be made before a hard frost. Make more than one application of this product if necessary to control re-growth of underground plant parts and seeds.

<b>Lotus, American</b>	2	0.75
------------------------	---	------

Apply this product when most target plants are at or beyond the bloom stage of growth. Enhanced results can be achieved when application is made during summer or fall months. Fall application must be made before a hard frost. Make more than one application of this product if necessary to control re-growth of underground plant parts and seeds.

<b>Melaleuca</b>	3	0.75
------------------	---	------

More than one application of this product will be needed for control, especially for vegetation partially submerged in water. Allow plants to re-grow to the 7- to 10-leaf stage before making next application.

<b>Milkweed, common</b>	2.3	1.5
<b>Milky, waterhen</b>	1.5 - 2.3	0.75

Apply this product when most target plants have reached the late-boot to flower stage of growth.

<b>Melick, common</b>	2.3 - 3.75	1.5
<b>Hopgrass</b>	2.3 - 3.75	1.5

Make application when most target plants are at least 6 inches in height (3- to 4-leaf stage of development) and actively growing.

PERENNIAL WEEDS RATE TABLE		
Weed Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
<b>Highbush blueberry, silverleaf</b>	2.3 - 3.75	1.5
Apply 3 to 3.75 quarts of this product per acre as a broadcast application west of the Mississippi River and 2.3 to 3 quarts per acre east of the Mississippi River when most target plants are at or beyond full bloom. Enhanced results can be obtained when application is made in late summer or fall after berries have formed.		
<b>Kentucky bluegrass, yellow</b>	2.3	0.75
Apply this product to control existing outside plants and attached immature nurseries when target plants are in flower or when new nurseries can be found at rhizome tips. Nurseries that have not germinated will not be controlled and will require repeated application of this product for long-term control.		
<b>Oxeye daisy</b>	1.5 - 2.3	0.75
Make application when most target plants have reached the boot to head stage of development. When applied prior to the boot stage, less than desirable control could be obtained in the fall; make application before plants have turned brown.		
<b>Panicum grass</b>	2.3 - 3.75	1.5
<b>Para grass</b>	3	0.75
More than one application of this product will be needed to achieve complete control. Allow plants to re-grow to the 1- to 10-leaf stage before making next application.		
<b>Perennial ryegrass</b>	3	1.5
<b>Phragmites*</b>	2 - 3.75	0.75 - 1.5
For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico, apply 3.75 quarts of this product per acre as a broadcast application or a 1.5-percent solution using a handheld sprayer. In other areas of the U.S., apply 2 to 3 quarts per acre as a broadcast application or for partial control, apply a 0.75-percent solution using a handheld sprayer. For enhanced results, make application in late summer or fall when plants are actively growing and in full bloom. Due to the dense nature of this vegetation (which can prevent good spray coverage) and uneven stages of growth, more than one application of this product might be necessary to achieve control. Visual symptoms of control will be slow to develop.		
<b>Quackgrass</b>	1.5 - 2.3	0.75
Apply this product when most target plants are at least 9 inches in height (3- to 5-leaf stage of development) and actively growing.		
<b>Rudbeckia*</b>	1.5	1.5
<b>Seed, common, giant</b>	3 - 3.75	1.5
For enhanced results make application in late summer or fall. Control can also be achieved by injecting 5 milliliters of this concentrated product (undiluted) directly into the second or third internode using a handheld application device. No surfactant is required.		
<b>Ryegrass, perennial</b>	1.5 - 2.3	0.75
Apply this product when most target plants have reached the boot to head stage of growth. When applied prior to the boot stage, enhanced control can result in the fall; make application before regrowth begins in spring.		

PERENNIAL WEEDS RATE TABLE		
Weed Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
<b>Sabine, giant</b>	3 - 3.75	2
Apply a 2-percent solution of this product containing 0.5 to 2 percent by volume of a nonionic surfactant approved for aquatic use and containing at least 70 percent active ingredient using spray-to-wet technique. For broadcast application, apply 3 to 3.75 quarts of this product per acre in 3 to 40 gallons of spray solution containing 0.1 percent by volume nonionic surfactant and 0.25 percent nonionic spreader-sticker. Surfactant approved for aquatic use. Allow a minimum of 3 days after application before disturbing vegetation in the application site. This product will not control plants that are completely submerged or have a majority of foliage under water.		
<b>Smartweed, swamp</b>	2.3 - 3.75	1.5
<b>Spotted dock</b>	3	0.75
Make application when most target plants are in full bloom. For enhanced results, apply in the summer or fall.		
<b>Spora, leafy*</b>	-	1.5
<b>Starrgrass, yellow</b>	-	1.5
<b>Sweet potato, wild*</b>	-	1.5
Make application when most target plants are at or beyond the bloom stage of growth. More than one application will be needed to achieve control.		
<b>Thistle, arctic*</b>	1.5 - 2.3	2
Make application when target plants are at or beyond the boot stage of growth.		
<b>Thistle, Canada</b>	1.5 - 2.3	1.5
Make application when target plants are at or beyond the boot stage of growth. Control can also be achieved by stem-injection. Cut 8 to 9 of tallest plants in a clump at bud stage. Push a tawny needle into the stem center and then slowly remove it as you inject 0.5 milliliter of this concentrated product into the stem. No surfactant required.		
<b>Timothy</b>	1.5 - 2.3	1.5
Make application when most target plants have reached the boot to head stage of development. If application is made prior to the boot stage, reduced control can result in the fall; make application before plants turn brown.		
<b>Tortoisegrass*</b>	3 - 3.75	0.75 - 1.5
Apply this product at a lower rate or spray solution concentration within the specified range when tortoisegrass is growing on terrestrial sites and at a higher rate or concentration within the range when partially submerged under water or growing as a floating mat. Additional applications of this product will be needed to maintain control.		
<b>Trumpetcrupper*</b>	1.5 - 2.3	1.5
<b>Ulex, common</b>	-	1.5
Make application to target plants at or beyond the seedhead stage of development. Visual symptoms will be slow to appear and might not appear for 3 or more weeks after application.		
<b>Vasegrass</b>	2.3 - 3.75	1.5
<b>Vetivergrass</b>	2.3 - 3.75	1.5

PERENNIAL WEEDS RATE TABLE		
Weed Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
<b>Waterhyacinth</b>	2.5 - 3	0.75 - 1
Make application when target plants are at or beyond the early bloom stage of development. Visual symptoms might require 3 or more weeks after application to appear, with complete necrosis and decomposition not occurring until 60 to 90 days after application. Use a higher application rate within the given range when more rapid visual effects are desired.		
<b>Waterhemp</b>	-	0.75 - 1
Apply a 1-percent solution of this product in areas of heavy infestation. Enhanced results can be obtained when applied from mid-summer through winter. Application in the spring could require more than one application to achieve control.		
<b>Waterprimrose</b>	-	0.75
Make application to target plants that are at or beyond the bloom stage of growth, but before fall color changes occur. Thorough coverage is necessary for enhanced control.		
<b>Wheatgrass, western</b>	1.5 - 2.3	0.75
Make application when most target plants have reached the boot to head stage of development. Application made prior to the boot stage could result in reduced control. In the fall, make application before plants turn brown.		

The application rate or spray solution concentration of the tank-mix product towards the higher end of the range. Refer to the individual product labels for approved uses and application rates. Not all tank-mix products listed are labeled for aquatic use.

**Aerial Aerial Herbicide Applicators Concentrate, Escort AP, Forestry Gator 4 Specialty, Forestry Gator XRT Specialty, Gator 3A Specialty, Gator 4 Specialty, Gator 4 Ultra Specialty, Vestal Specialty emulsifier, methoxy, methoxy**

Ensure that the proper amount of the Gator herbicide is thoroughly mixed with water in the spray tank before adding this product.

#### Cut Stump Application

This product may be used to control re-growth and re-sprouting of woody brush and tree on any site listed on this label. On the woody brush or tree close to the soil or water surface and immediately apply a 50 to 100 percent (undiluted) solution of this product to the freshly cut surface using an applicator capable of applying this product to the entire cut surface. A delay in application could result in reduced performance. For enhanced control on the woody brush or tree during period of active growth and full leaf expansion and apply this product. No surfactant is needed for cut stump application.

For control of the tree of heaven (*Ailanthus altissima*), cut the tree above to the soil surface and immediately apply a 50 percent solution of this product (10 fluid ounces per quart of solution) and 10 percent Aerial Herbicide L3 to 4 fluid ounces per quart of solution) in water to the freshly cut surface. **DO NOT MAKE A CUT STUMP APPLICATION WHEN THE ROOTS OF DESIRABLE WOODY BRUSH OR TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP AS INJURY COULD OCCUR IN THE ADJACENT TREES.** Some species stems or trees can share a common root system. Adjacent trees having a similar age, height and spacing could be an indication of a shared root system. Whether grafted or shared, injury of likely to occur to adjacent stems or trees when this product is applied to one or more trees sharing common root system.

#### Woody Brush and Tree Injection and Frill Application

This product may be used to control woody brush and trees listed in this section by injection or frill application on any aquatic and terrestrial site listed on this label.

Inject or apply the equivalent of 1 milliliter (0.04 fluid ounce) of this product for every 2 to 3 inches of trunk diameter at breast height (DBH). If injecting this product into the woody brush or tree, use equipment capable of penetrating into the living plant tissue under the bark. No surfactant is required for direct injection of this product into woody brush and trees.

For frill application, apply a 50 to 95-percent solution of this product in water with 0.5% or more by volume of a nonionic surfactant. To achieve continuous frill around the tree or to cut evenly spaced around the tree between all branches. As tree diameter increases, enhanced results can be achieved by applying this product to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow this product to run out of the minor cut areas in species that freely exude sap. Make the frill or cut at an oblique angle to produce a "weeping effect" and apply a 95-percent solution of this product with a nonionic surfactant as described above. For enhanced results, make this application during periods of active growth and after full leaf expansion.

#### Modified High-Volume and Low-Volume Backpack Application

For control and partial control of woody brush, trees and vines listed on this label when using a backpack sprayer or other handheld equipment and a directed tree volume foliar application technique, apply a 4 to 6 percent solution of this product containing 0.5 to 1 percent by volume of a nonionic surfactant evenly over the plant crown to cover 50 percent of the upper foliage of undesirable woody brush, trees and vines.

WOODY BRUSH, TREES AND VINES RATE TABLE

Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
Alder	2.3 - 3	0.75 - 1.2
Ash*	1.5 - 3.75	0.75 - 1.5
Aspen, quaking	1.5 - 2.3	0.75 - 1.2
Barelyever (Bourmar)*	1.5 - 3.75	0.75 - 1.5
Beech*	1.5 - 3.75	0.75 - 1.5
Birch	1.5	0.75
Blackberry	2.3 - 3	0.75 - 1.2
Blackgum	1.5 - 3.75	0.75 - 1.5
Bracon	1.5 - 3.75	0.75 - 1.5
Broom, French, Scotch	1.5 - 3.75	1.2 - 1.5
Buckhorn, California*	1.5 - 3	0.75 - 1.5
Casahuate	1.5 - 3.75	0.75 - 1.5
Caster bean	1.5 - 3.75	1.5

Also for control of 4 milliliters of this concentrated (undiluted) product per quart directly into the lower portion of the main stem using a handheld injection device. No surfactant is required.

For partial control, apply this product when at least 50 percent of the new leaves are fully developed.

Ceanothus*	1.5 - 3.75	0.75 - 1.5
Cherry, bitter, black, pin	1.5 - 3.75	1 - 1.5
Cottonwood, eastern	1.5 - 3.75	0.75 - 1.5
Coyote brush	2.3 - 3	1.2 - 1.5

WOODY BRUSH, TREES AND VINES RATE TABLE

Species	Broadcast Rate (quarts/acre)	Handheld Spray-to-Wet Concentration (% solution)
Kudzu	3	1.5
Locust, black*	1.5 - 3	0.75 - 1.5
Madrone, red*	-	1.5
Magnolia, sweetbay	1.5 - 3.75	0.75 - 1.5
Mazzeutia*	1.5 - 3.75	0.75 - 1.5
Maple, red	1 - 3.75	0.75 - 1.2

For control, apply a 0.75- to 1.2 percent solution of this product using a handheld sprayer when leaves are fully developed. For partial control, apply 1 to 3 quarts per acre as a broadcast application.

For control, apply this product using a handheld sprayer when at least 50 percent of the new leaves are fully developed.

Maple, sugar	0.75 - 1.2	
For control, apply this product using a handheld sprayer when at least 50 percent of the new leaves are fully developed.		
Maple, vine*	1.5 - 3.75	0.75 - 1.5
Monkey flower*	1.5 - 3	0.75 - 1.5
Oak, black, white*	1.5 - 3	0.75 - 1.5
Oak, northern pin	1.5 - 3	0.75 - 1.2
For control, apply this product when at least 50 percent of the new leaves are fully developed.		
Oak, poison	3 - 3.75	1.5
Repeat applications might be required to maintain control. Application in the test must be made before leaves lose green color.		
Oak, post	2.3 - 3	0.75 - 1.2
Oak, red	-	0.75 - 1.2

For control, apply this product using a handheld sprayer when at least 50 percent of the new leaves are fully developed.

Oak, scrub*	1.5 - 3	0.75 - 1.5
Oak, southern red	1.5 - 3.75	1 - 1.5
Orange, Osage	1.5 - 3.75	0.75 - 1.5
Peppercorn, Brazilian (Florida holly)*	1.5 - 3.75	1.5
Periwinkle*	1.5 - 3.75	0.75 - 1.5
Pine	1.5 - 3.75	0.75 - 1.5
Poplar, yellow*	1.5 - 3.75	1 - 1.5
Prunus	2.3 - 3	0.75 - 1.2
Rasperry	1.5 - 3.75	0.75 - 1.5
Rhodod, eastern	1.5 - 3.75	0.75 - 1.5
Rhodod, western	1.5 - 3.75	0.75 - 1.5
Rose, multiflora	1.5	0.75
Make application prior to leaf feeding insects		
Ruscus, white*	1.5 - 3.75	0.75 - 1.5
Sage, black	1.5 - 3	0.75
Sage, white*	1.5 - 3	0.75 - 1.5
Sapindus, California	1.5 - 3	0.75
Salmonberry	1.5	0.75
Satibush	-	1



**WOODY BRUSH, TREES AND VINES RATE TABLE**

Species	Broadcast Rate (quarts per acre)	Handheld Spray-In- Wick Concentration (% active)
Saltcedar*	3 - 3.75	1 - 2
For partial control, apply a 1- to 2-percent solution of this product using a handheld sprayer or 3 to 3.75 quarts per acre as a broadcast application.		
For control, apply a 1- to 1.5-percent solution of this product in a tank-mix with Arsenal Herbicide or Arsenal Herbicide Application Concentrate using a handheld sprayer. For control using broadcast application, apply 1.5 quarts of this product per acre in a tank-mix with an appropriate rate of Arsenal Herbicide or Arsenal Herbicide Application Concentrate to plants less than 6 feet tall. To control saltcedar greater than 6 feet tall using broadcast application, apply 3 quarts of this product per acre on a tank-mix with a higher rate of Arsenal Herbicide or Arsenal Herbicide Application Concentrate.		
Sassaparil*	1.5 - 3.75	0.75 - 1.5
Sea Hurtle	1	1
Sorrel*	1.5 - 3.75	0.75 - 1.5
Sumac: laurel, poison, smooth, sugarbush, winged*	1.5 - 3	0.75 - 1.5
Sweetgum	1.5 - 2.3	0.75 - 1.5
Swordfern*	1.5 - 3.75	0.75 - 1.5
Tailwren, Chinese	0.75	0.75
Tan oak re-sprouts*	1.5	1.5
Thimbleberry	1.5	0.75
Tobacco, tree*	1.5 - 3	0.75 - 1.5
Toyon*	1.5	1.5
Trumpetcrusher	1.5 - 2.3	0.75 - 1.2
Vine maple*	1.5 - 3.75	0.75 - 1.5
Virginia creeper	1.5 - 3.75	0.75 - 1.5
Wasmuth's, southern*	1.5 - 3.75	1.5
Willow	2.3	0.75
Yerba Santa, California*	1.5	1.5

**Other woody brush and trees listed on this label:** For partial control, apply 1.5 to 3.75 quarts of this product per acre as a broadcast application or a 0.75- to 1.5-percent solution using a handheld sprayer and a spray-to-wet application technique.

**130 LIMIT OF WARRANTY AND LIABILITY**

Bayer CropScience LP ("Company") warrants that this product conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Bayer and all users shall use this product only for the purposes of and in accordance with the Complete Directions for Use label ("Directions") and shall expressly notify this Company of any claim, whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, Bayer and all users are responsible for all tests, injuries or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, crop injury or failure of this product to control weeds or types which develop resistance to glyphosate, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, use and/or application in any manner not explicitly set forth in or inconsistent with the Directions, moisture conditions outside the moisture range specified in the Directions or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product (except in accordance with this Company's stewardship requirements and with express written permission from this Company).

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE, SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT RECEIVED OR AN ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLENISHMENT OF SUCH QUANTITY OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. NO EXCEPT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, Bayer and all users are deemed to have accepted the terms of this Limit of Warranty and Liability, which may not be revised by any verbal or written agreement. If terms are not acceptable, return at once unopened.

In case of an emergency involving this product, or for medical assistance call collect, day or night, 1-800-334-7577

Roundup, Roundup Custom and Design, and TRUEBLUE ADVANTAGE PROVEN RELIABLE, SUPPORTED and Design, are trademarks of Bayer Group. All other trademarks are the property of their respective owners. ©2020 Bayer Group. All rights reserved.

EPA Reg. No. 524-343

Packed for  
BAYER CROPSOURCE LP  
800 N. LINDBERGH BLVD  
ST. LOUIS, MISSOURI 63167 U.S.A.

# Roundup CUSTOM<sup>®</sup>

**FOR AQUATIC & TERRESTRIAL USE**



Un herbicida de amplio espectro para aplicación postemergencia, para el control de malezas acuática e industrial, ornamental, en céspedes, forestación, lados de carreteras, servidumbres de paso, cultivos selectos y otros usos terrestres indicados.

(Para una lista completa de aplicaciones acuáticas y terrestres, vea la sección Modo de empleo de esta etiqueta).

**INGREDIENTE ACTIVO:**

\*Glicosato, N-(fosfonometil) glicina en la forma de su sal de isopropilamina 53.8%  
 OTROS INGREDIENTES: 46.2%  
 100.0%

\* Contiene 648 gramos por litro o 5.4 libras por galón norteamericano del ingrediente activo glicosato, en la forma de su sal de isopropilamina, lo cual es equivalente a 480 gramos por litro o 4.0 libras por galón norteamericano (39.9% por peso) del ácido glicosato.

Nº. Reg. EPA 524-343 EPA Est. 524-IA-1

Manténgase Fuera del Alcance de los Niños  
**PRECAUCIÓN**

Consulte el interior para conocer las precauciones adicionales.  
**INSTRUCCIONES DE USO COMPLETAS**

No todos los productos recomendados en esta etiqueta han sido registrados para su uso en California. Verifique el estado de registro de cada producto en California antes de utilizarlo.

**PARA INFORMACIÓN SOBRE EL PRODUCTO O AYUDA PARA UTILIZAR ESTE PRODUCTO, LLAME GRATIS AL 1-866-99BAYER (1-866-992-2937)**

**EN CASO DE EMERGENCIA RELACIONADA CON ESTE PRODUCTO O PARA AYUDA MEDICA, LLAME POR COBRAR, DE DIA O DE NOCHE, AL 1-800-334-7577**

Envasado para:  
 BAYER CROPSCIENCE LP  
 800 N. LINDBERGH BLVD.  
 ST. LOUIS, MISSOURI 63167 USA  
 ©2020 Grupo Bayer. Todos los derechos reservados.

lea toda la etiqueta antes de usar este producto  
 Use solo de acuerdo con las instrucciones de la etiqueta  
 Lea la declaración "LÍMITE DE GARANTÍA Y RESPONSABILIDAD" al final de esta etiqueta antes de comprar o usar. Si los términos no son aceptables devuélvalos de inmediato sin abrir.  
 EVITE EL CONTACTO DE ESTE HERBICIDA CON FOLIAJE, TALLOS VERDES, FRUTOS NO LIGEROS, EXPUESTAS O FRUTAS DE CULTIVOS, PLANTAS Y ARBOLCS DESEABLES. COMO PODRÍA PRODUCIRSE UNA LESIÓN GRAVE A LA PLANTA O LA DESTRUCCIÓN.  
 ESTE ES UN PRODUCTO DE USO FINAL. BAYER CROPSOURCE LP NO TIENE LA INTENCIÓN Y NO LO HA REGISTRADO PARA SU REFORMULACIÓN O REDISEÑO.

**CONTENIDO**

1 1.0 INGREDIENTES 18

2 2.0 NÚMEROS DE TELÉFONO IMPORTANTES 18

3 3.0 DECLARACIONES PREVENTIVAS 18

3.1 Riesgos para los seres humanos y los animales domésticos 18

3.2 Riesgos para el medio ambiente 18

3.3 Riesgos físicos o químicos 18

4 4.0 ALMACENAMIENTO Y ELIMINACIÓN 19

5 5.0 INFORMACIÓN DEL PRODUCTO 19

6 6.0 MANEJO DE RESISTENCIA DE MALEZAS 19

6.1 Prácticas de manejo de malezas 19

6.2 Manejo de biotipos resistentes al glifosato 19

7 7.0 MEZCLAS 20

7.1 Mezclar con agua 20

7.2 Sulfonilamida 20

7.3 Mezclas de tanque 20

7.4 Procedimiento de mezcla en tanque 20

7.5 Mezclar concentraciones de solución de rocio 20

7.6 Colorantes y tintes 21

7.7 Aditivos de reducción de dispersión 21

8 8.0 EQUIPOS Y TÉCNICAS PARA LA APLICACIÓN 21

8.1 Manejo de la dispersión del rocio 21

8.2 Equipo de aplicación aérea 21

8.3 Equipo de aplicación terrestre 22

8.4 Bovederos manuales 22

8.5 Equipo de aplicación selectiva 22

8.6 Sistemas por inyección 23

8.7 Aplicación por goteo controlado (GDA) 23

9 9.0 SITIOS DE USO ACUÁTICO Y TERRESTRE 23

9.1 Sitios acuáticos 23

9.2 Sitios terrestres 24

10 10.0 INFORMACIÓN ADICIONAL SOBRE MANEJO DEL LUGAR 24

10.1 Manejo de bosques, árboles de madera y árboles de Navidad 24

10.2 Manejo de baldíos de vida silvestre y especies nativas 25

10.3 Manejo de vivero ornamental y de producción 25

10.4 Manejo de áreas con cultivos (vegetables y hortalizas) 25

10.5 Manejo de zonas de pasturas 25

10.6 Manejo de fertilizantes 25

10.7 Manejo de tierras de pastoreo 26

10.8 Manejo de bidos de carreteras 26

10.9 Manejo de servicios públicos 26

11 11.0 USOS EN CULTIVOS 26

11.1 Cultivos en árboles, enredaderas y arbustos 26

11.1.1 Cultivos de frutas cítricas 27

11.2 Cultivos anuales y perennes 27

11.2.1 Caña de azúcar 28

11.3 Producción de semillas de pasto y forrajes 29

12 12.0 MALEZAS CONTROLADAS 29

12.1 Control de malezas: remoción y manejo químico en especies 29

12.2 Malezas anuales 30

12.3 Malezas perennes 30

12.4 Arbustos, enredaderas y matorrales leñosos 33

13 13.0 LÍMITES EN LA GARANTÍA Y LA RESPONSABILIDAD 35

**1.0 INGREDIENTES**

**INGREDIENTE ACTIVO**

\*Glifosato, (1-(4-isopropilfenilo) glicolato, 53.8%

en la forma de su sal de isopropilamina 46.2%

**OTROS INGREDIENTES**

\*Contiene 640 gramos por litro o 5.4 libras por galón (normalmente) del ingrediente activo glifosato, en la forma de su sal de isopropilamina. lo cual es equivalente a 480 gramos por litro o 4.0 libras por galón (normalmente) (33.9% por peso) del acido glifosico

**2.0 NÚMEROS DE TELÉFONO IMPORTANTES**

1 PARA INFORMACIÓN SOBRE EL PRODUCTO: LLAME AL NÚMERO DE EMERGENCIAS PARA AYUDAR ESTE PRODUCTO. LLAME GRATIS AL 1-800-952-4373

2 EN CASO DE EMERGENCIA RELACIONADA CON ESTE PRODUCTO PARA AYUDA MÉDICA: LLAME POR COPIAR EL NÚMERO DE EMERGENCIAS AL 1-800-324-1977

**3.0 DECLARACIONES PREVENTIVAS**

**3.1 Riesgos para los seres humanos y los animales domésticos**

**Manténgase Fuera del Alcance de los Niños**

**PRECAUCIÓN**

**ANIMALES DOMÉSTICOS** Se considera que este producto es relativamente no tóxico para perros y otros animales domésticos. Sin embargo, la ingestión de este producto o de abundantes cantidades de vegetación (cualquiera fuere) podría causar náuseas, gastrointestinales temporales (vómitos, diarrea, calambos, etc.). Si se presentan dichos síntomas, de al animal suficiente cantidad de líquidos para evitar la deshidratación. Llame a un veterinario si los síntomas persisten por más de 24 horas.

**Equipo de protección personal (EPP)**

Los usuarios y personas que manipulan este producto deben usar, como mínimo mangas largas y pantalones largos, zapatos y rebatacos.

Respeche las instrucciones del fabricante para limpiar y mantener los equipos de protección personal (EPP). En caso de que no haya instrucciones, use detergente y agua caliente. Mantenga el EPP aparte del resto de la ropa y lávelo por separado.

**Recomendaciones de seguridad para el usuario:**

Los usuarios deben:

- lavarse las manos antes de comer, beber, masticar chicle, usar tabaco o usar el baño
- Quitar la ropa de inmediato si el pesticida les respasa la ropa. Luego deben lavarse muy bien y ponerse ropa limpia

**3.2 Riesgos para el medio ambiente**

Matar las malezas acuáticas puede dar lugar a una reducción o pérdida de oxígeno en el agua debido a la descomposición de material vegetal muerto. Esta pérdida de oxígeno puede causar que los peces se asfixien. Antes de aplicar pesticidas a aguas naturales consulte con la agencia de su estado que es la principal responsable de su regulación para determinar si se necesita un permiso. Para usos terrestres, no aplique directamente al agua, en áreas donde el agua superficial este presente o en áreas intermareales por debajo del nivel medio de mareas altas, excepto si se aplica por aire por encima de la cobertura forestal. No contamine el agua cuando limpie el equipo o deslacte el agua de lavado y empaque del equipo.

**3.3 Riesgos físicos o químicos**

Para mezclar, almacenar y aplicar la solución de rocio de este producto se pueden usar recipientes de acero inoxidable, fibra de vidrio, plástico o recipientes de acero recubiertos internamente con plástico.

**NO MEZCLE, ALMACENE O APlique ESTE PRODUCTO O LAS SOLUCIONES DE ROCIO DE ESTE PRODUCTO EN EMBAJES DE ACERO GALVANIZADO O EN REVESTIMIENTO (EXCEPTO ACERO INOXIDABLE) O EN TANQUES DE ROCIO.** Si se utiliza en estos embaes o tanques, este producto o las soluciones de rocio de este producto reaccionan y producen gas hidrógeno que puede formar una mezcla de gases altamente inflamable. Esta mezcla de gases podría encenderse o explotar si está en contacto con fuego, chispas, sopletes para soldar, respiradores encendidos o cualquier otra fuente de ignición y causar lesiones personales graves.

**INSTRUCCIONES DE USO**

Se considera una violación a la ley federal usar este producto de una manera que no sea la indicada en la etiqueta. Este producto solo puede utilizarse de acuerdo con las instrucciones de uso en la etiqueta o según las etiquetas complementarias que se publican por separado. Puede solicitar las etiquetas complementarias para este producto a su vendedor minorista autorizado de Bayer CropScience LP, o a su representante de Bayer CropScience LP.

No aplique este producto de manera que entre en contacto con los trabajadores u otras personas, no sea directamente o por arastre. Solamente las aplicaciones que usen protección pueden estar en el área durante su aplicación. Para verificar requisitos expuestos de su título o estado, consulte con la agencia responsable de la regulación del uso de pesticidas.

**Requisitos para uso agrícola**

Utilice este producto solo de acuerdo con la etiqueta y con las Normas de Protección para Trabajadores, 40 CFR Parte 170. Estas normas contienen los requisitos para la protección de trabajadores agrícolas en granjas, bosques, viveros e invernaderos y para las personas que manipulan pesticidas agrícolas. Contienen los requisitos para capacitar, desarrollar, notificar y ofrecer asistencia de emergencia. También contienen instrucciones específicas y excepciones relativas a las alarmaciones en esta etiqueta sobre los equipos de protección personal (EPP), los refugios de acceso restringido. Los requisitos en esta caja se referirán únicamente a las aplicaciones de este producto cubiertas por las Normas de Protección para Trabajadores.

No entre ni permita la entrada de personal a los áreas tratadas durante el intervalo de entrada restringida (IER) por sus reglas en inglés) de 3 horas. El EPP que se requiere para el acceso anticipado a zonas tratadas de acuerdo con las Normas de Protección para Trabajadores y que incluye el contacto con material tratado, como plantas, tierra o agua es: overoles, zapatos con cañutines y guantes resistentes a sustancias químicas confeccionados con cualquier tipo de material impermeable.

**Requisitos para usos no agrícolas**

Los requisitos en esta caja se refieren a las aplicaciones de este producto que NO cubren las Normas de Protección para Trabajadores para pesticidas agrícolas (40 CFR Parte 170). Las Normas se aplican cuando este producto se utiliza para producir plantas agrícolas en granjas, bosques, viveros o invernaderos.

Manténgase a las personas y a las mascotas fuera de las áreas tratadas hasta que la solución de rocio se haya secado.

















Control de matorrales, arbustos y enredaderas

Este producto se puede usar para controlar árboles, enredaderas y matorrales...

HEZOLA DE JANDUE: Este producto se puede aplicar en una mezcla de...

Arrestal, Herbicida contactante para aplicaciones: Arsenal, Chopper, Clapper...

Control de malezas en Bermudagrass latente y en crecimiento activo

Este producto se puede usar para controlar (a) o parcialmente controlar...

107 Manejo de tierras de pastoreo

Este producto controla o inhibe muchas malezas anuales que crecen en...

Para controlar la invasión de malezas de pastos anuales en tierras de pastoreo...

Control de Bromera: Una aplicación al voleo de 9 a 12 onzas líquidas de este...

Control de Hierbabuena: Para controlar o inhibir las plantas de hierbabuena...

quemado. Repite la aplicación anualmente para eliminar las semillas de...

108 Manejo de lodos de carreteras

Todos los usos de este producto descritos en esta etiqueta pueden utilizarse...

HEZOLA DE JANDUE: Este producto puede mezclarse en tanque con los...

Altra 11, Altra 12, Altra 13, Altra 14, Altra 15, Altra 16, Altra 17, Altra 18, Altra 19...

109 Manejo de servicios públicos

Este producto se puede usar a lo largo de arboledas de paso de...

HEZOLA DE JANDUE: Este producto se puede mezclar en tanque con los...

Altra 11, Altra 12, Altra 13, Altra 14, Altra 15, Altra 16, Altra 17, Altra 18, Altra 19...

Aguerre de que el producto (a) no es bien mezclado con agua de...

Para obtener mejores resultados con el recorte lateral, aplique este producto...

110 USOS EN CULTIVOS

11.1 Cultivos de árboles, enredaderas y arbustos

ESTA SECCIÓN OFECE INSTRUCCIONES DE USO QUE APPLICAN A TODOS LOS...

INSTRUCCIONES DE USO: A menos que se prohiba específicamente en...

Aplique 12 onzas líquidas a 4 cuartos de galón de este producto por acre...













**TABLA DE PROPORCIONES PARA MALEZAS PERENNES**

Especies de maleza	Proporción por dilución (cuartos de galón/ha)	Concentración del roció para mejor resultado de mano (% solución)
Guinea grass	2.3	0.75

Aplique este producto cuando la mayoría de las plantas objetivo hayan alcanzado la etapa de crecimiento de 7 hojas.

**Hemlock, poison**  
También se puede lograr el control inyectando 5 mililitros de inyección manual en una cana de una hoja por planta. 17 pulgadas por encima de la corona de la raíz. No se requiere surfactante.

**Hogweed, giant**  
Inyecte 5 mililitros de una solución al 5 por ciento de este producto en una lirreite 5 milímetros por encima de la corona de la raíz. No se requiere surfactante.

**Horseweed**  
2.3 - 3.75 1.5  
No se requiere surfactante.

**Horseweed**  
3 1.5  
Aplique este producto cuando la mayoría de las plantas objetivo hayan alcanzado la etapa de desarrollo de brotación tardía a floración. Para obtener mejores resultados, aplique a finales del verano o en otoño.

**Horsetail, field**  
Inyecte 0.5 mililitros de este producto por tallo directamente en el tallo de la planta, un segmento por encima de la corona de la raíz. No se requiere surfactante.

**Isoplat**  
1.5 1.5

**Iris, yellow flag**  
Corte los tallos de flores 8 o 9 pulgadas por encima de la corona de la raíz. Introdúzca una aguja inyectora en el centro del tallo y luego retirela lentamente a medida que inyecta 0.5 mililitros de este producto con un inyector manual. No se requiere surfactante.

**Ivy, cape, German**  
1.5 - 3 0.75 - 1.5

**Juncus/lemn artificialis**  
2.3 - 3.75 1.5

**Johnsongrass**  
1.5 - 2.3 0.75

Aplique este producto cuando la mayoría de las plantas objetivo hayan alcanzado la etapa de desarrollo de bola a final de verano o antes de que las plantas se hayan oscurecido en otoño. Cuando se aplica antes de la etapa de bola, el control puede verse reducido.

**Knapweed**  
1.5 - 2.3 0.75

**Knapweed**  
3 1.5

Aplique este producto cuando la mayoría de las plantas objetivo hayan alcanzado la etapa de desarrollo de brotación tardía a floración. Para obtener mejores resultados, aplique a finales del verano o en otoño.

**TABLA DE PROPORCIONES PARA MALEZAS PERENNES**

Especies de maleza	Proporción por dilución (cuartos de galón/ha)	Concentración del roció para mejor resultado de mano (% solución)
Knapweed, Bohemian, giant, Japanese	3	2

Aplique 3 cuartos de galón de este producto por acre como aplicación al voleo en 3 a 40 galones de solución de roció con 0.5 a 1 por ciento por volumen de un surfactante no iónico. Para aplicar usando un recodo de roció y la técnica de roció para mejor, aplique una solución al 2 por ciento de este producto que contiene de 0.5 a 2 por ciento por volumen de un surfactante no iónico. Para obtener mejor control, no altere la repetición en el área de aplicación por un mínimo de 7 días después de la aplicación.

También se puede obtener control al hacer un corte limpio de los tallos justo por debajo del segundo o tercer nodo sobre la superficie y aplicar 0.36 onzas líquidas (10 mililitros) de una solución al 50 por ciento de este producto en agua en el poco o riego que quede. Asegúrese de que los restos de la parte superior de la planta que se eliminó se recojan y desechen correctamente para evitar que se propaguen nuevas plantas de los retoños.

El uso de bobarras, tales como paneles de cartón, maderas o cajas hechas de plástico, ayudará a reducir la propagación de materia vegetal. La proporción de aplicar un total combinado de este producto no puede exceder 8 cuartos de galón por acre. También se puede lograr el control inyectando 5 mililitros de este producto por tallo en el segundo o tercer internodo usando un dispositivo de inyección manual. No se requiere surfactante.

**Lantana**  
0.75 - 1

Aplique cuando la mayoría de las plantas se encuentren en la etapa de floración o después de ella. Mida la concentración más alta de solución de roció para plantas que han alcanzado la etapa de crecimiento leñoso.

**Lespedeza**  
2.3 - 3.75 1.5

**Loosestrife, purple**  
2 1 - 1.5

Aplique cuando la mayoría de las plantas se encuentren en la etapa de floración o después de ella. Se pueden obtener mejores resultados cuando la aplicación se realiza durante los meses de verano o de otoño. La aplicación en otoño debe hacerse antes de una helada agresiva.

**Lupinus, American**  
2 0.75

Aplique cuando la mayoría de las plantas se encuentren en la etapa de floración o después de ella. Se pueden obtener mejores resultados cuando la aplicación se realiza durante los meses de verano o de otoño. La aplicación en otoño debe hacerse antes de una helada agresiva. Puede ser necesaria más de una aplicación de este producto para controlar el crecimiento de semillas y partes de plantas subterráneas.

**Melaleuca**  
3 0.75

Se necesitan más de una aplicación de este producto para lograr el control. Especialmente para la vegetación parcialmente sumergida en el agua. Permita que las plantas crezcan hasta la etapa de 7 a 10 hojas antes de realizar la siguiente aplicación.

**Milkwed, common**  
2.3 1.5

Aplique este producto cuando la mayoría de las plantas objetivo hayan alcanzado la etapa de desarrollo de brotación tardía a floración.

**Minty, wirestem**  
1.5 - 2.3 0.75

Aplique cuando la mayoría de las plantas objetivo tengan por lo menos 8 pulgadas de alto (etapa de desarrollo de 3 a 4 hojas) y estén en crecimiento activo.

**Mullein, common**  
2.3 - 3.75 1.5

**Nasopirgrass**  
2.3 - 3.75 1.5

**TABLA DE PROPORCIONES PARA MALEZAS PERENNES**

Especies de maleza	Proporción por dilución (cuartos de galón/ha)	Concentración del roció para mejor resultado de mano (% solución)
Hipocistis, silverleaf	2.3 - 3.75	1.5

Aplique de 2 a 3.75 cuartos de galón de este producto por acre como aplicación al voleo al oeste del río Mississippi y de 2 a 3 cuartos de galón por acre al este del río Mississippi cuando la mayoría de las plantas objetivo estén en plena floración o después. Se pueden obtener mejores resultados cuando la aplicación se realiza a finales del verano o en otoño, después de la formación de tallos.

**Horseweed, purple, amarillo**  
2.3 0.75

Aplique este producto para controlar las plantas existentes de horseweed y las nuevas plantas que surgen cuando las plantas objetivo están en flor o cuando se pueden encontrar nuevas plantas en las purdas de los rizomas. No se pueden controlar las plantas que todavía no han germinado y será necesario repetir las aplicaciones de este producto para lograr el control a largo plazo.

**Orchardgrass**  
1.5 - 2.3 0.75

Aplique cuando la mayoría de las plantas objetivo hayan alcanzado la etapa de desarrollo de bola a floración. Cuando se aplica antes de la etapa de bola, puede obtenerse un menor control del desarrollo. En otoño, aplique el tratamiento antes de que las plantas se oscurezcan.

**Pennis grass**  
2.3 - 3.75 1.5

**Pennis grass**  
3 0.75

Se necesitan más de una aplicación de este producto para lograr el control total. Permita que las plantas crezcan hasta la etapa de 7 a 10 hojas antes de realizar la siguiente aplicación.

**Peppercorn, purple**  
3 1.5

**Perennial\***  
2 - 3.75 0.75 - 1.5

Para el control manual de piragüas en Florida y en los condados de otros estados que bordean el Golfo de México, aplique 3.75 cuartos de galón de este producto por acre como aplicación al voleo o una solución al 1.5 por ciento usando un rociador de mano. En otros áreas de los EE. UU., aplique de 2 a 3 cuartos de galón por acre como aplicación al voleo o para control manual, aplique una solución al 0.75 por ciento usando un rociador de mano.

Para obtener mejores resultados, aplique a finales del verano o en otoño cuando las plantas estén en crecimiento activo y en plena floración. Déjado a la naturaleza densa de la vegetación que puede impedir la correcta cobertura del roció) y a las etapas de crecimiento irregulares, puede ser necesaria más de una aplicación de este producto para lograr el control. Los síntomas visuales de control se desarrollarán con lentitud.

**Quackgrass**  
1.5 - 2.3 0.75

Aplique este producto cuando la mayoría de las plantas objetivo tengan por lo menos 8 pulgadas de alto (etapa de desarrollo de 3 a 4 hojas) y estén en crecimiento activo.

**Rhynchos\***  
1.5 1.5

**River, common, pigweed**  
3 - 3.75 1.5

Para obtener mejores resultados, aplique a finales del verano o en otoño. También se puede lograr el control inyectando 5 mililitros de este producto concentrado (1 un galón) directamente en el segundo o tercer internodo usando un dispositivo de inyección manual. No se requiere surfactante.

**Ryegrass, perennal**  
1.5 - 2.3 0.75

Aplique este producto cuando la mayoría de las plantas objetivo hayan alcanzado la etapa de crecimiento de bola a brotación. Cuando se aplica antes de la etapa de bola, el control puede verse reducido. En el otoño aplique el tratamiento antes de que el invierno se oscurezca.

**TABLA DE PROPORCIONES PARA MALEZAS PERENNES**

Especies de maleza	Proporción por dilución (cuantos de recipientes de 1 galón)	Concentración del rocio para mejor en recipientes de 1 galón	Concentración del rocio para mejor en recipientes de 1 galón (% solución)
Sambino, plant	3 - 3.75	2	0.75 - 1

Aplique una solución al 2 por ciento de este producto que contenga de 0.5 a 2 por ciento por volumen de un surfactante no iónico aprobado para uso acuático y por volumen de 10 por ciento de ingrediente activo usando la técnica de rocio para maso. Para aplicación al viento, aplique de 3 a 3.75 cuantos de galón de este producto por acre en 3 a 40 galones de solución de agua que contenga 0.1 por ciento por volumen de ingrediente activo y 0.75 por ciento de surfactante no iónico o mezcla de rocio coproporcionado y 0.75 por ciento de surfactante no iónico. Deje transcurrir al menos 3 días después de la aplicación antes de alterar la vegetación en el área de aplicación. Este producto no controla plantas que están completamente sumergidas o que tienen la mayoría de su follaje debajo del agua.

Smartweed, Swamp	2.3 - 3.75	1.5	0.75
Spartan	3	0.75	0.75

Aplique cuando la mayoría de las plantas objetivo estén en plena floración. Para obtener mejores resultados, aplique en verano o en otoño.

Spartan, hairy*	1.5 - 2.3	1.5	1.5
Starbistle, amarillo	1.5 - 2.3	1.5	1.5
Sweet potato, wild*	1.5	1.5	1.5

Aplique cuando la mayoría de las plantas objetivo se encuentren en la etapa de floración o después de ella. Puede ser necesario más de una aplicación para el control total.

Thistle, arctic	1.5 - 2.3	2	1.5
Thistle, Canada	1.5 - 2.3	1.5	1.5

Aplique cuando la mayoría de las plantas objetivo se encuentren en la etapa de floración o después de ella.

Timothy	1.5 - 2.3	1.5	1.5
---------	-----------	-----	-----

Aplique cuando la mayoría de las plantas objetivo hayan alcanzado la etapa de desarrollo de baya a floración. Si la aplicación se hace antes de la etapa de baya, el control puede verse reducido. En el otoño, aplique el tratamiento antes de que las plantas se oscurezcan.

Trifolium	3 - 3.75	0.75 - 1.5	0.75 - 1.5
-----------	----------	------------	------------

Aplique este producto en una proporción o concentración de solución de rocio menor dentro del rango especificado cuando los productos se creen en el campo y en una proporción o concentración mayor dentro del rango cuando está parcialmente sumergido en agua o crecido como mata flotante. Se necesitan aplicaciones adicionales de este producto para mantener el control.

Trifolium	1.5 - 2.3	1.5	1.5
Taraxacum	1.5 - 2.3	1.5	1.5
Vasegrass	2.3 - 3.75	1.5	1.5
Wetgrass	2.3 - 3.75	1.5	1.5

Aplique a las plantas objetivo cuando se encuentren en la etapa de floración o después de ella. Los síntomas visuales aparecerán lentamente y puede que no aparezcan hasta 3 semanas o más después de la aplicación.

**TABLA DE PROPORCIONES PARA MALEZAS PERENNES**

Especies de maleza	Proporción por dilución (cuantos de recipientes de 1 galón)	Concentración del rocio para mejor en recipientes de 1 galón	Concentración del rocio para mejor en recipientes de 1 galón (% solución)
Waterhyacinth	2.5 - 3	0.75 - 1	0.75 - 1

Aplique cuando las plantas objetivo se encuentren en la etapa de floración temprana o después de ella. Los síntomas visuales pueden tardar en aparecer 3 o más semanas después de la aplicación y la necrosis completa y descomposición de 80 a 90 días después de la aplicación. Si desea obtener los efectos visuales más rápido, use una proporción de aplicación más alta dentro del rango indicado.

Waterhyacinth	2.5 - 3	0.75 - 1	0.75 - 1
---------------	---------	----------	----------

Aplique una solución al 1 por ciento de este producto en áreas de floración severa. Se pueden obtener mejores resultados si se aplica de mediados del verano hasta el invierno. Se aplica en primavera es posible que se necesite más de una aplicación para lograr el control.

Waterhyacinth	1.5 - 2.3	0.75	0.75
---------------	-----------	------	------

Resque la aplicación a plantas objetivo que se encuentran en etapa de floración o después de ella, pero antes de que cambien de color en el otoño. Para obtener un mejor control es necesaria una cobertura completa.

Waterhyacinth	1.5 - 2.3	0.75	0.75
---------------	-----------	------	------

Aplique cuando la mayoría de las plantas objetivo hayan alcanzado la etapa de desarrollo de baya a floración. Las aplicaciones realizadas antes de la etapa de baya pueden tener menor control. En el otoño, aplique el tratamiento antes de que las plantas se oscurezcan.

Waterhyacinth	1.5 - 2.3	0.75	0.75
---------------	-----------	------	------

Aplique cuando la mayoría de las plantas objetivo se encuentren en la etapa de floración o después de ella. Puede ser necesario más de una aplicación para el control total.

Waterhyacinth	1.5 - 2.3	0.75	0.75
---------------	-----------	------	------

Aplique cuando la mayoría de las plantas objetivo se encuentren en la etapa de floración o después de ella. Puede ser necesario más de una aplicación para el control total.

Waterhyacinth	1.5 - 2.3	0.75	0.75
---------------	-----------	------	------

Aplique cuando la mayoría de las plantas objetivo se encuentren en la etapa de floración o después de ella. Puede ser necesario más de una aplicación para el control total.

enredaderas y matorrales leñosos del sitio de la aplicación. Pueden ser necesarias aplicaciones adicionales de este producto para controlar subterráneos.

**MEZCLAS DE TANQUE:** Este producto puede aplicarse en cualquier proporción indicada en esta etiqueta en una mezcla de tanque con los siguientes productos para aumentar el espectro de control de malezas herbáceas, árboles, enredaderas y matorrales leñosos. Para el control de las malezas herbáceas, aplique el producto de la mezcla de tanque más baja dentro del rango especificado. Para el control de poblaciones densas o de árboles, enredaderas y matorrales leñosos difíciles de controlar, aumente la proporción de aplicación o la concentración de solución de rocio del producto en la mezcla de tanque hacia el extremo más alto del rango. Consulte las etiquetas de cada producto para conocer los usos aprobados y las proporciones de aplicación. No todos los productos en mezcla de tanque están aprobados para uso acuático.

Arsenal - Herbicida concentrado para aplicaciones Aerial, Escort XP, Forestry, Galfon 4, Specialty, Forestry, Galfon XR, Specialty, Galfon 3A, Specialty, Galfon 4 Specialty, Galfon 4 Ultra Specialty, Vastlan Specialty, amazepr, metsulfuron methyl, triclopyr.

Asegúrese de que la cantidad correcta del herbicida Galfon está bien mezclada con agua en el tanque de rocio antes de agregar este producto a la aplicación.

Este producto se puede usar para controlar rebrotos y retoños de matorrales leñosos y árboles en cualquier sitio indicado en esta etiqueta. Corte el árbol o matorral leñoso cerca de la superficie de la tierra o del agua y aplique inmediatamente una solución de este producto al 50 o 100 por ciento (con diluir) a la superficie recién cortada utilizando un pulverizador capaz de aplicar este producto a todo el cambium. La demora en la aplicación puede resultar en un rendimiento inferior. Para obtener mejores resultados, corte el árbol o matorral leñoso durante el período de crecimiento activo y expánsion completa de los liños y aplique este producto. No se necesita surfactante para la aplicación a troncos cortados.

Para controlar lires de hoaces (Maidenhair tree), corte el árbol cerca de la superficie de la tierra y aplique de inmediato una solución al 50 por ciento de este producto (15 onzas líquidas por cuarto de galón de solución) y un 10 por ciento de herbicida Arsenal (de 1 a 3 onzas líquidas por cuarto de galón de solución en agua) a la superficie recién cortada.

NO REALICE UNA APLICACIÓN EN TRONCOS CORTADOS CUANDO LAS RAÍCES DE MATORRALES LEÑOSOS O ARBOLES DECADENTES PUEDAN ESTAR INFLUENCIADAS EN LAS RAÍCES DEL TONCO CORTADO PORQUE SI PUEDEN CAUSAR DAÑO A LOS ARBOLES ADYACENTES. Algunos retoños de árboles pueden compartir el mismo sistema de raíces. Los árboles adyacentes de edad, altura y espaciamiento similares pueden tener raíces compartidas. Ya sea injertados o conectados, es probable que se dañen y árboles adyacentes cuando se aplica este producto a uno o más árboles que comparten un sistema común de raíces.

Aplique de Inyección y anillado de matorrales leñosos y árboles. Este producto se puede usar para controlar árboles y matorrales leñosos indicados en esta sección por aplicación de inyección o anillado (fill) en cualquier sitio acuático o terrestre indicado en esta etiqueta. Inerte o aplique el equivalente 1 mililitro (0.01 onzas líquidas) de este producto por cada 2.0 pulgadas de diámetro del tronco a la altura del pecho (DBH en inglés). Si aplica este producto en un matorral leñoso o árbol, use el mayor nivel de penetración (el mayor nivel de la placa debajo de la corteza). No se requiere surfactante para la inyección directa de este producto en árboles y matorrales leñosos.

Para aplicación con anillado, aplique una solución de 50 a 95 por ciento de este producto en agua con 0.5% o más por volumen de un surfactante no iónico a una incisión ambar continua alrededor del árbol o en cortes espaciados uniformemente alrededor del árbol por debajo del nivel de las ramas. A medida que el diámetro del árbol aumenta, pueden obtenerse mejores resultados al aplicar este producto a una incisión ambar continua o en cortes menos espaciados. Evite las técnicas de aplicación que permitan

que este producto se escuma de las áreas cortadas o con incisiones. En las especies que se sequegan se recomienda hacer las cortes o incisiones en un ángulo oblicuo para producir un efecto de coque y aplicar una solución al 5% por ciento de este producto con un surfactante no iónico, como se describe a continuación. Para obtener mejores resultados, realice esta aplicación durante el período de crecimiento activo y después de la expansión completa de las hojas.

**Aplicación modificada de alto volumen y bajo volumen con mochila**

Para el control total y parcial de áfidos, arañitas y matorrales leñosos indicado en esta etiqueta, al usar un rociador de mochila u otro equipo manual y una técnica de aplicación foliar original de bajo volumen, aplique una solución de 4 a 8 por ciento de este producto que contenga de 0.5 a 1 por ciento por volumen de un surfactante no iónico de forma uniforme sobre la corona de la planta para cubrir el 50 por ciento del follaje superior de los árboles, enredaderas y matorrales leñosos no deseados.

**TABLA DE PROPORCIONES PARA MATORRALES LEÑOSOS, ÁRBOLES Y ENREDADERAS**

Especie	Proporción por dilución (cuantos de galón/acre)	Concentración del rociador para mejorar en rociador de mano (% solución)
Alder	2.3 - 3	0.75 - 1.2
Asit*	1.5 - 3.75	0.75 - 1.5
Aspen, quaking	1.5 - 2.3	0.75 - 1.2
Bearbrier (Bearmat)*	1.5 - 3.75	0.75 - 1.5
Beech*	1.5 - 3.75	0.75 - 1.5
Birch	1.5	0.75
Blackberry	2.3 - 3	0.75 - 1.2
Blackgum	1.5 - 3.75	0.75 - 1.5
Bracken	1.5 - 3.75	0.75 - 1.5
Broom: French, Scotch	1.5 - 3.75	1.2 - 1.5
Buckwheat, California*	1.5 - 3	0.75 - 1.5
Casahuate*	1.5 - 3.75	0.75 - 1.5
Castor bean	1.5 - 3.75	1.5
Además, para controlar, injecte 4 mililitros de este producto concentrado (un drácula) por planta directamente en la parte inferior del tallo principal con un dispositivo de inyección manual. No requiere surfactante.		
Catsclaw*		1.2 - 1.5
Para controlar parcial, aplique este producto cuando por lo menos el 50 por ciento de las hojas nuevas estén completamente desarrolladas.		
Deamollas*	1.5 - 3.75	0.75 - 1.5
Chamisa*	1.5 - 3.75	0.75
Ceanoth: amarga, negro, rojo	1.5 - 3.75	1 - 1.5
Cottonwood, eastern	1.5 - 3.75	0.75 - 1.5
Coyote brush	2.3 - 3	1.2 - 1.5
Para controlar, aplique este producto cuando por lo menos el 50 por ciento de las hojas nuevas estén completamente desarrolladas.		
Cypress: swamp, bald	1.5 - 3.75	0.75 - 1.5
Deerweed	1.5 - 3.75	0.75 - 1.5
Dewberry	2.3 - 3	0.75 - 1.2
Dogwood*	3 - 3.75	1 - 2
Elderberry	1.5	0.75
Elm*	1.5 - 3.75	0.75 - 1.5
Encarnillo, blue gem		1.5
Para controlar nuevos brotes de entalpias, aplique este producto usando un rociador manual cuando los nuevos brotes tengan entre 6 a 12 pies de altura. Asegúrese de conseguir una cobertura completa.		

**TABLA DE PROPORCIONES PARA MATORRALES LEÑOSOS, ÁRBOLES Y ENREDADERAS**

Especie	Proporción por dilución (cuantos de galón/acre)	Concentración del rociador para mejorar en rociador de mano (% solución)
Gallberry	1.5 - 3.75	0.75 - 1.5
Gorse*	1.5 - 3.75	0.75 - 1.5
Hackberry, western	1.5 - 3.75	0.75 - 1.5
Haystack*	1.5 - 3	0.75 - 1.5
Hawthorn	1.5 - 2.3	0.75 - 1.2
Hazel	1.5	0.75
Hickory*	3 - 3.75	1 - 2
Honeyuckle	2.3 - 3	0.75 - 1.2
Hornbeam, American*	1.5 - 3.75	0.75 - 1.5
Huckleberry	1.5 - 3.75	0.75 - 1.5
Ivy, poison	3 - 3.75	1.5
Kudzu	3	1.5
Locust, black*	1.5 - 3	0.75 - 1.5
Madrone (strawberry)		1.5
Magnolia, sweetbay	1.5 - 3.75	0.75 - 1.5
Marrubium*	1.5 - 3.75	0.75 - 1.5
Maple, red	1 - 3.75	0.75 - 1.2
Para controlar, aplique una solución de 0.75 a 1.2 por ciento de este producto usando un rociador manual cuando las hojas estén completamente desarrolladas. Para control parcial, aplique de 1 a 3.75 cuartos de galón por acre en aplicaciones al voleo.		
Maple, sugar		0.75 - 1.2
Para controlar, aplique este producto con un rociador manual cuando por lo menos el 50 por ciento de las hojas nuevas estén completamente desarrolladas.		
Maple, vine*	1.5 - 3.75	0.75 - 1.5
Monkey flower*	1.5 - 3	0.75 - 1.5
Oak, black, white*	1.5 - 3	0.75 - 1.5
Oak, northern, pin	1.5 - 3	0.75 - 1.2
Para controlar, aplique este producto cuando por lo menos el 50 por ciento de las hojas nuevas estén completamente desarrolladas.		
Oak, poison	3 - 3.75	1.5
Para controlar, aplique este producto para lograr el control. La aplicación en otoño debe hacerse antes de que las hojas pierdan su color verde.		
Oak, post	2.3 - 3	0.75 - 1.2
Oak, red		0.75 - 1.2
Para controlar, aplique este producto con un rociador manual cuando por lo menos el 50 por ciento de las hojas nuevas estén completamente desarrolladas.		
Oak, scrub*	1.5 - 3	0.75 - 1.5
Oak, southern red	1.5 - 3.75	1 - 1.5
Orange, Osage	1.5 - 3.75	0.75 - 1.5
Peppercorn, Brazilian (Florida holly)*	1.5 - 3.75	1.5
Persimmon*	1.5 - 3.75	0.75 - 1.5
Pine	1.5 - 3.75	0.75 - 1.5
Poplar, amantilla*	1.5 - 3.75	0.75 - 1.5
Prunus	1.5 - 3.75	1 - 1.5

**TABLA DE PROPORCIONES PARA MATORRALES LEÑOSOS, ÁRBOLES Y ENREDADERAS**

Especie	Proporción por dilución (cuantos de galón/acre)	Concentración del rociador para mejorar en rociador de mano (% solución)
Raspberry	2.3 - 3	0.75 - 1.2
Rubus, eastern	1.5 - 3.75	0.75 - 1.5
Rubus, western	1.5 - 3.75	0.75 - 1.5
Rose, multiflora	1.5	0.75
Realice las aplicaciones antes de que los insectos que se alimentan de las hojas estén en las hojas.		
Russian olive*	1.5 - 3.75	0.75 - 1.5
Sage, black	1.5 - 3	0.75
Sage, white*	1.5 - 3	0.75 - 1.5
Sagebrush, California	1.5 - 3	0.75
Salmonberry	1.5	0.75
Sambucus		1
Sambucus*	3 - 3.75	1 - 2
Para el control parcial, aplique una solución de 1 a 2 por ciento de este producto usando un rociador manual o de 3 a 3.75 cuartos de galón por acre en aplicación al voleo. Para controlar, aplique una solución de 1 a 1.5 por ciento de este producto en una mezcla de lanque con herbicida Arsenal o Herbicida concentrado para aplicadores Arsenal usando un rociador manual. Para controlar con aplicación al voleo, aplique 1.5 cuartos de galón de este producto por acre en una mezcla de lanque con una proporción más alta de herbicida Arsenal o Herbicido concentrado para aplicadores Arsenal.		
Sassafras*	1.5 - 3.75	0.75 - 1.5
Sea Myrtle		1
Sourwood*	1.5 - 3.75	0.75 - 1.5
Sumac: laurel, poison, smooth, sugarbush, winged*	1.5 - 3	0.75 - 1.5
Sweetgum	1.5 - 2.3	0.75 - 1.5
Swordfern*	1.5 - 3.75	0.75 - 1.5
Tallowtree, Chinese		0.75
Tan oak (triflorata)*	1.5	1.5
Thimbleberry	1.5	0.75
Tobacco, lres*	1.5 - 3	0.75 - 1.5
Typha*		1.5
Trumpetcreeper	1.5 - 2.3	0.75 - 1.2
Vine maple*	1.5 - 3.75	0.75 - 1.5
Virginia creeper	1.5 - 3.75	0.75 - 1.5
Waxmyrtle, southern*	1.5 - 3.75	1.5
Willow	2.3	0.75
Yerba Santa, California*		1.5

\* Control parcial  
Otros árboles y matorrales leñosos indicados en esta etiqueta - Para control parcial, aplique de 1.5 a 3.75 cuartos de galón de este producto por acre como aplicación al voleo o una solución al 0.75 o 1.5 por ciento usando un rociador manual y la técnica de rocio para mosca.

### 13.0

## LIMITES EN LA GARANTIA Y LA RESPONSABILIDAD

Bayar Corporation IP (la "Compañía") garantiza que este producto conforma con la descripción, apariencia de la etiqueta, HASTA EL GRADO QUE SEA COMPATIBLE CON LA LEGISLACIÓN PERTINENTE. NO SE HACE NINGUNA OTRA GARANTIA EXPRESA O IMPLICITA, AGUNDA DE LA IDONEIDAD PARA UN USO PARTICULAR O COMERCIALIZACIÓN. Esta garantía está sujeta también a las condiciones y limitaciones que aquí se indican.

El comprador y todos los usuarios utilizarán este producto únicamente con los propósitos y de acuerdo con la etiqueta de las Instrucciones completas para el uso ("Instrucciones") y notificarán de inmediato a la Compañía si hacen alguna reclamación que se base en un contrato, negligencia, estricta responsabilidad u otros derechos estatutarios.

Hasta el grado que sea compatible con la legislación pertinente, el comprador y todos los usuarios son responsables por todos los daños, lesiones o daños que resultasen por el uso o manipulación en condiciones que estén más allá del control de esta Compañía, incluyendo sus límites de uso, incompatibilidad con productos que no sean los señalados en las Instrucciones, aplicación o contacto con vegetación que no se cuera destruir, caída a caminos o la incapacidad del producto para controlar los tipos de malezas que de otro modo no estarían controladas. Condiciones climáticas adversas, condiciones climáticas fuera de las normas que se consideren normales en el lugar de la aplicación y para el período de tiempo en el cual se aplica el producto, así como condiciones climáticas que estén fuera de los límites indicados en las Instrucciones, uso y/o aplicación que no estén específicamente aconsejadas en las Instrucciones o no sean compatibles con estas, condiciones de humedad que estén fuera de los límites establecidos en las Instrucciones, o la presencia de productos en la tierra o sobre ella en los cultivos o en la vegetación que se está tratando, diferencias a los indicados en las Instrucciones.

Esta Compañía no garantiza ninguno de los productos reformulados o reempacados de este producto, excepto de acuerdo a los requisitos de la administración de esta Compañía y con el permiso escrito expreso de esta Compañía.

HASTA EL GRADO QUE SEA COMPATIBLE CON LA LEGISLACIÓN PERTINENTE, LA ÚNICA EXCLUSIVA COMPENSACIÓN AL USUARIO O COMPRADOR Y EL LÍMITE DE RESPONSABILIDAD DE ESTA COMPAÑÍA O DE CUALQUIER OTRO VENDEDOR POR CUALQUIER PERDIDA O POR TODAS LAS FERMIDAS, FERMENTOS O DAÑOS QUE RESULTEN DEL USO O MANEJO DE ESTE PRODUCTO INCLUYENDO RECLAMOS QUE SE BASAN EN UN CONTRATO, NEGLIGENCE, ESTRUCTA RESPONSABILIDAD U OTROS DERECHOS EXTRACONTRACTUALES, SERÁ EL PRECIO PAGADO POR EL USUARIO O EL COMPRADOR POR LA CANTIDAD INVOLUCRADA EN ESTE PRODUCTO O A ELECCIÓN DE ESTA COMPAÑÍA O DE OTRO VENDEDOR, EL REEMPLAZO DE DICHA CANTIDAD O SI NO SE OBLIGÓ ANTES DE LA COMPRA SE REEMPLAZARÁ DICHA CANTIDAD DEL PRODUCTO HASTA EL GRADO QUE SEA COMPATIBLE CON LA LEGISLACIÓN PERTINENTE. EN NINGUN CASO ESTA COMPAÑÍA U OTRO VENDEDOR, SERÁN RESPONSABLES POR DAÑOS INDIRECTALES, CONSECUENTES O ESPECIALES.

En el momento de abrir y usar el producto, se asume que el comprador y todos los usuarios han aceptado las condiciones de los LÍMITES EN LA GARANTIA Y LA RESPONSABILIDAD que no pueden darse por medio de ningún acuerdo verbal o escrito. Si las condiciones son inaceptables, devuelve el producto inmediatamente sin abrir el empaque.

En caso de emergencia relacionada con este producto o si necesita ayuda médica, llame por teléfono de día o de noche, al 1-800-334-7577.

Certainty, Durrider, Roundup, Roundup Condon y su diseño, y TRIEBLUE ADVANTAGE PROVEN RELIABLE SUPPORTED y su diseño, son marcas registradas de Grupo Bayer. Todas las demás marcas registradas son propiedad de sus respectivos dueños. © 2020 Grupo Bayer. Todos los derechos reservados.

Nº Reg. EPA 522-343

Entregado por:  
BAYER CORPORATION IP  
500 N. LINDBERGH BLVD  
ST LOUIS, MISSOURI 63167 U.S.A

20  
CODE

Roundup® CUSTOM

## FOR AQUATIC & TERRESTRIAL USE

A broad-spectrum postemergence herbicide for aquatic and industrial, turf, ornamental, forestry, roadside, utility rights-of-way, select crop, and other listed terrestrial weed control.

(For a complete list of aquatic and terrestrial use sites, see the Directions for Use section of the attached labeling.)

### ACTIVE INGREDIENT:

\*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt..... 53.8%

OTHER INGREDIENTS:..... 46.2%  
100.0%

\*Contains 648 grams of the active ingredient glyphosate in the form of its isopropylamine salt per liter, or 5.4 pounds per U.S. gallon, which is equivalent to 480 grams of the acid glyphosate, per liter or 4.0 pounds per U.S. gallon (39.9% by weight)

EPA Reg. No. 524-343

EPA Est. 524-1A-1

## Keep Out of Reach of Children

## CAUTION

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Not all products listed on this label are registered for use in California.

Check the registration status of each product in California before using.

See Complete Directions for Use attached to this label for complete Agricultural and Non-Agricultural Use Requirements of the Worker Protection Standard, Directions for Use and Limit of Warranty and Liability in English and Spanish.

GROUP

9

HERBICIDE

## STORAGE AND DISPOSAL

Proper pesticide storage and disposal are essential to protect against exposure to people and the environment due to leaks and spills, excess product or waste, and vandalism. Do not allow this product to contaminate water, foodstuffs, feed or seed by storage and disposal.

**PESTICIDE STORAGE:** STORE ABOVE 5 F (-15°C) TO KEEP PRODUCT FROM CRYSTALLIZING. Crystals will settle to the bottom. If allowed to crystallize, warm to 68 F (20°C) to redissolve and roll or shake container or recirculate contents of larger containers to mix well before using. Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep containers closed to prevent spills and contamination. See individual container label for additional storage conditions. If any

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in the container, including rinsate, by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program. Such programs are often run by state or local governments or by industry. All disposal must be in accordance with applicable federal, state and local regulations and procedures.

**CONTAINER HANDLING AND DISPOSAL:** Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in this container. Contact your state regulatory agency to determine allowable practices in your state. Triple rinse or pressure rinse (or equivalent) this container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Continue to drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Place container so that it can drain directly into application equipment or mix tank while rinsing, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle into the side of the container and rinse at about 40 PSI for at least 30 seconds. Continue to drain for 10 seconds after the flow begins to drip. Once properly rinsed, some plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest collection site, contact your chemical dealer or Bayer CropScience LP at 1-866-99BAYER (1-866-992-2937). If recycling is not available, dispose of in accordance with federal, state and local regulations and procedures, which may include puncturing the properly rinsed container and disposing in a sanitary landfill.

FOR PRODUCT INFORMATION OR ASSISTANCE USING THIS PRODUCT,  
CALL TOLL-FREE, 1-866-99BAYER (1-866-992-2937)

IN CASE OF AN EMERGENCY INVOLVING THIS PRODUCT OR FOR MEDICAL  
ASSISTANCE, CALL COLLECT, DAY OR NIGHT, 1-800-334-7577

Roundup and Roundup Custom and Design are trademarks of Bayer Group.

© 2020 Bayer Group. All rights reserved.

Packed for:

BAYER CROPSCIENCE LP  
800 N. LINDBERGH BLVD.  
ST. LOUIS, MISSOURI 63167 USA



US62217691B

1910189v2 10/20

NET 2.5 GAL

883580926201

PLACE BAR CODE HERE

